

Gelfand–Zeitlin theory from the perspective of classical mechanics II

BERTRAM KOSTANT* and NOLAN WALLACH**

ABSTRACT. In this paper, Part II, of a two part paper we apply the results of [KW], Part I, to establish, with an explicit dual coordinate system, a commutative analogue of the Gelfand-Kirillov theorem for $M(n)$, the algebra of $n \times n$ complex matrices. The function field $F(n)$ of $M(n)$ has a natural Poisson structure and an exact analogue would be to show that $F(n)$ is isomorphic to the function field of a $n(n-1)$ -dimensional phase space over a Poisson central rational function field in n variables. Instead we show that this the case for a Galois extension, $F(n, \mathfrak{e})$, of $F(n)$. The techniques uses a maximal Poisson commutative algebra of functions arising from Gelfand-Zeitlin theory, the algebraic action of a $n(n-1)/2$ -dimensional torus on $F(n, \mathfrak{e})$, and the structure of a Zariski open subset of $M(n)$ as a $n(n-1)/2$ -dimensional torus bundle over a $n(n+1)/2$ -dimensional base space of Hessenberg matrices.

0. Part II continuation of Introduction

0.6. We recall some of the notation and results in Part I, i.e., [KW]. If k is a positive integer then $I_k = \{1, \dots, k\}$. $M(n)$ is the algebra of all $n \times n$ complex matrices. If $m \in I_n$, then regard $M(m) \subset M(n)$ as the upper left block of all $m \times m$ matrices. If $x \in M(n)$ then $x_m \in M(m)$ is the upper left principal $m \times m$ minor of x . Using a natural isomorphism of (the Lie algebra) $M(n)$ with its dual space, $M(n)$ becomes a Poisson manifold so that its affine ring $\mathcal{O}(M(n))$ is a Poisson algebra. For any $k \in \mathbb{Z}_+$ let $d(k) = k(k+1)/2$. The subalgebra $J(n)$ of $\mathcal{O}(M(n))$, generated by the symmetric polynomial $Gl(m)$ -invariants of $M(m)$ for all $m \in I_n$, is a polynomial algebra with $d(n)$ generators and more importantly it is a maximal Poisson commutative subalgebra of $\mathcal{O}(M(n))$.

In Part I we showed that the Poisson vector field ξ_f on $M(n)$ corresponding to any $f \in J(n)$ is globally integrable on $M(n)$ and a choice of generators of $J(n)$ defines an abelian Lie group A of dimension $d(n-1)$ operating on $M(n)$. The orbits of A are explicitly determined in Part I and the orbits are independent of the choice of generators. One particular choice are the functions, $p_i(x)$, $i \in I_{d(n)}$, $x \in M(n)$,

* Research supported in part by NSF grant DMS-0209473 and in part by the KG&G Foundation

** Research supported in part by NSF grant MTH 0200305

where, for all $m \in I_n$, $p_{d(m-1)+k}(x)$, $k \in I_m$, are the nontrivial coefficients of the characteristic polynomial of x_m .

A suitable measure on $\mathbb{R}$ and the Gram-Schmidt process define a sequence, $\phi_k(t)$, $k \in \mathbb{Z}_+$, of orthogonal polynomials on $\mathbb{R}$. Let W_n be the span of ϕ_{m-1} , $m \in I_n$, and let $x \in M(n)$ be the matrix, with respect to this basis, of the operator of multiplication by t , followed by projection on W_n . The matrix x is Jacobi and for $m \in I_n$ one recovers the orthogonal polynomial ϕ_m as the characteristic polynomial of x_m . In particular the all important zeros of the orthogonal polynomials ϕ_m appear as the eigenvalues of the x_m . One motivation for our work here is to set up Poisson machinery to deal with the eigenvalues of x_m for any $x \in M(n)$. In the course of setting up this machinery we have obtained a number of new results. Some of these results have appeared in Part I ([KW]). In the present paper, Part II, we will be concerned with establishing a refinement of a commutative analogue of the Gelfand-Kirillov theorem. The refinement refers to exhibiting an explicit coordinate system satisfying the Poisson commutation relations of phase space. The coordinate system emerges from the action of an algebraic group and the structure of $M(n)$, obtained in Part I, as sort of a cotangent bundle over the variety of Hessenberg matrices.

In more detail let $M_\Omega(n)$ be the Zariski open (dense) subset of $M(n)$ defined as the set of all $x \in M(n)$ such that x_m is regular semisimple in $M(m)$ for all $m \in I_n$ and such that the spectrum of x_{m-1} , for $m > 0$, has empty intersection with the spectrum of x_m . In Part I $M_\Omega(n)$ was shown to have the following structure: It is a $(\mathbb{C}^\times)^{d(n-1)}$ bundle over a $d(n)$ -dimensional base space ($d(n-1) + d(n) = n^2$) base space. The fibers are not only the level sets for the functions in $J(n)$ but also the fibers are the orbits of A in $M_\Omega(n)$. The base space, denoted by $\mathfrak{b}_{e,\Omega(n)}$, is the intersection $\mathfrak{b}_e \cap M_\Omega(n)$ where $\mathfrak{b}_e$ is the space of Hessenberg matrices. That is, $x \in \mathfrak{b}_e$ if and only if x is of the form

$$x = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n-1} & a_{1n} \\ 1 & a_{22} & \cdots & a_{2n-1} & a_{2n} \\ 0 & 1 & \cdots & a_{3n-1} & a_{3n} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & a_{nn} \end{pmatrix}$$

Also $\mathfrak{b}_{e,\Omega(n)}$ is Zariski dense in $\mathfrak{b}_e$. Theorem 2.5 in Part I concerning establishing a beautiful property of $\mathfrak{b}_e$ plays a major role here. In part II a “Lagrangian” property of $\mathfrak{b}_e$ plays a key role in showing the dual coordinates s_i , defined below, Poisson commute. Part I has 3 sections. A serious deficiency in $M_\Omega(n)$ in dealing with the commutative analogue of the Gelfand-Kirillov theorem is that one cannot consistently solve the characteristic polynomials of x_m , for all $m \in I_n$ and all $x \in M_\Omega(n)$ to yield algebraic eigenvalue functions r_i on $M_\Omega(n)$.

0.7. We begin, in the first section of Part II, labeled Section 4, to obtain such functions on a covering space $M_\Omega(n, \mathfrak{e})$ of $M_\Omega(n)$. Initially the covering map

$$\pi_n : M_\Omega(n, \mathfrak{e}) \rightarrow M_\Omega(n)$$

is only understood to be analytic. The covering admits, as deck transformations, a group, Σ_n , isomorphic to the direct product of the symmetric groups S_m , $m \in I_n$ and as analytic manifolds

$$M_\Omega(n, \mathfrak{e})/\Sigma_n \cong M_\Omega(n)$$

However much more structure is needed and established in §4. For one thing $M_\Omega(n, \mathfrak{e})$ is a nonsingular affine variety and π_n is a finite étale morphism. For another, if $F(n)$ is the field of rational functions on $M(n)$ and $F(n, \mathfrak{e})$ is the field of rational functions on $M_\Omega(n, \mathfrak{e})$, then $F(n, \mathfrak{e})$ is a Galois extension of $F(n)$ with Σ_n as Galois group. Furthermore the affine ring $\mathcal{O}(M_\Omega(n, \mathfrak{e}))$ is the integral closure of $\mathcal{O}(M_\Omega)$ in $F(n, \mathfrak{e})$. Very significant for our purposes, there exists (eigenvalue) functions $r_i \in \mathcal{O}(M_\Omega(n, \mathfrak{e}))$, $i \in I_{d(n)}$ with the property that for any $m \in I_n$ and any $z \in M_\Omega(n, \mathfrak{e})$ the numbers $r_{d(m-1)+k}(z)$, $k \in I_m$ are the eigenvalues of x_m , where $x = \pi_n(z)$. The Poisson structure on $M_\Omega(n)$ lifts to $M_\Omega(n, \mathfrak{e})$ and one has $[r_i, r_j] = 0$ for all $i, j \in I_{d(n)}$.

0.8. §5 is devoted to the construction of the dual coordinates $s_j \in \mathcal{O}(M_\Omega(n, \mathfrak{e}))$, $j \in I_{d(n-1)}$. There are two key points here. (a) It is shown that the Poisson vector fields $\xi_{\mathfrak{r}_i}$ on $M_\Omega(n, \mathfrak{e})$ integrate and generate a complex algebraic torus, $A_{\mathfrak{r}} \cong (\mathbb{C}^\times)^{d(n-1)}$ which operates algebraically on $M_\Omega(n, \mathfrak{e})$ and in fact if $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ is the π_n inverse image of $\mathfrak{b}_{\mathfrak{e}, \Omega(n)}$ in $M_\Omega(n, \mathfrak{e})$, then the map

$$A_{\mathfrak{r}} \times M_\Omega(n, \mathfrak{e}, \mathfrak{b}) \rightarrow M_\Omega(n, \mathfrak{e}), \quad (b, y) \mapsto \mathfrak{b} \cdot y$$

is an algebraic isomorphism. The natural coordinate system on $A_{\mathfrak{r}}$ then carries over to $M_\Omega(n, \mathfrak{e})$ defining functions $s_j \in \mathcal{O}(M_\Omega(n, \mathfrak{e}))$, $j \in I_{d(n-1)}$, when they are normalized so that, for all j , s_j is the constant 1 on $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$. The second key point, (b), yields the Poisson commutativity $[s_i, s_j] = 0$ from the Lagrangian property of $\mathfrak{b}_e$. See Theorem 5.20 and its proof. Combining Theorems 5.14 and 5.23 one has

Theorem 0.16. *The image of the map*

$$M_\Omega(n, \mathfrak{e}) \rightarrow \mathbb{C}^{n^2}, \quad z \mapsto (r_1(z), \dots, r_{d(n)}(z), s_1(z), \dots, s_{d(n-1)}(z)) \quad (0.13)$$

is a Zariski open set Y in $\mathbb{C}^{n^2}$ and (0.13) is an algebraic isomorphism of $M_\Omega(n, \mathfrak{e})$ with Y . Furthermore one has the following Poisson commutation relations:

$$\begin{aligned} (1) \quad & [r_i, r_j] = 0, \quad i, j \in I_{d(n)} \\ (2) \quad & [r_i, s_j] = \delta_{ij} s_j, \quad i \in I_{d(n)}, \quad j \in I_{d(n-1)} \\ (3) \quad & [s_i, s_j] = 0, \quad i, j \in I_{d(n-1)} \end{aligned} \quad (0.14)$$

Noting that s_i vanishes nowhere on $M_\Omega(n, \mathfrak{e})$ one has $r_{(i)} \in \mathcal{O}(M_\Omega(n, \mathfrak{e}))$ for $i \in I_{d(n-1)}$ where $r_{(i)} = r_i/s_i$. Replacing r_i by $r_{(i)}$ in (2) one has the more familiar phase space commutation relation $[r_{(i)}, s_j] = \delta_{ij}$. For the implication of Theorem 0.16 on the structure of the field $F(n, \mathfrak{e})$ see Theorem 5.24.

0.9. Of course given the eigenvalue functions r_i , the dual coordinates s_j are not uniquely determined. In the present paper they are given by the use of the algebraic group A_τ (defined by the r_i) and a set of Hessenberg matrices as a base space. Independently and quite differently the papers [GKL1] and [GKL2] also deal with establishing a refined commutative analogue of the Gelfand-Kirillov theorem. A point of similarity is the use of the coordinates r_i and the necessity, thereby, to go to a covering. In §3 of [GKL1] dual coordinates are given, denoted in that paper by Q_{nj} . It seems to be an interesting question to write down equations expressing a relation between the Q_{nj} in [GKL1] and the s_i here. The first three sections of the paper are in Part I. Part II begins with §4.

4. The covering $M_\Omega(n, \mathfrak{e})$ of $M_\Omega(n)$ and the eigenvalue functions r_i

4.1. We retain the general notation of Part 1 so that n is a positive integer and $M(n)$ is the space of all complex $n \times n$ matrices. As in (2.61), for $m \in I_n$, (see §1.1) let $\mathfrak{d}(m)$ be the space of all diagonal matrices in $M(m)$ and let $\mathfrak{e}(m)$ be the (connected) Zariski open subset of all regular elements in $\mathfrak{d}(m)$. That is, if $z \in \mathfrak{d}(m)$, then $z \in \mathfrak{e}(m)$ if and only if the diagonal entries of z are distinct. Consider the direct product

$$\mathfrak{e} = \mathfrak{e}(1) \times \cdots \times \mathfrak{e}(m) \quad (4.1)$$

so that if $\nu \in \mathfrak{e}$ we can write

$$\nu = (\nu(1), \dots, \nu(n)) \quad (4.2)$$

where $\nu(m) \in \mathfrak{e}(m)$. In addition we will write

$$\nu(m) = \text{diag}(\nu_{1m}, \dots, \nu_{mm}) \quad (4.3)$$

where the numbers $\nu_{im} \in \mathbb{C}$, $i \in I_m$, are distinct. Taking notation from (2.53) let $\mathfrak{e}_{\Omega(n)}$ be the Zariski open subset of $\mathfrak{e}$ defined so that if $\nu \in \mathfrak{e}$ then $\nu \in \mathfrak{e}_{\Omega(n)}$ if and only if

$$\nu_{im} \neq \nu_{jm+1}, \quad \forall m \in I_{n-1}, i \in I_m, j \in I_{m+1} \quad (4.4)$$

Of course $\mathfrak{e}_{\Omega(n)}$ is a nonsingular variety where

$$\dim \mathfrak{e}_{\Omega(n)} = d(n)$$

(see §0.1). The symmetric group S_m , as the Weyl group of $(M(m), \mathfrak{d}(m))$, operates freely on $\mathfrak{e}(m)$ and the direct product $\Sigma_n = S_1 \times \cdots \times S_n$ (a group of order $\prod_{m \in I_n} m!$) operates freely on $\mathfrak{e}_{\Omega(n)}$ where if $\sigma = (\sigma_1, \dots, \sigma_n)$, $\sigma_m \in S_m$, is in Σ_n and $\nu \in \mathfrak{e}_{\Omega(n)}$ then, using the notation of (4.2) and (4.3),

$$\sigma \cdot \nu = (\sigma_1 \cdot \nu(1)), \dots, \sigma_m \cdot \nu(m) \quad (4.5)$$

and

$$\sigma_m \cdot \nu(m) = \text{diag}(\nu_{\sigma_m^{-1}(1)m}, \dots, \nu_{\sigma_m^{-1}(m)m}) \quad (4.6)$$

4.2. Recall the Zariski open set $M_{\Omega}(n)$ of $M(n)$ (see (2.53)). In particular we recall that the matrices in $M_{\Omega}(n)$ are regular semisimple. Consider the direct product $\mathfrak{e}_{\Omega(n)} \times M_{\Omega}(n)$ and let

$$M_{\Omega}(n, \mathfrak{e}) = \{(\nu, x) \in \mathfrak{e}_{\Omega(n)} \times M_{\Omega}(n) \mid \nu(m) \text{ is } Gl(m)\text{-conjugate to } x_m, \forall m \in I_n\} \quad (4.7)$$

It is clear that $M_{\Omega}(n, \mathfrak{e})$ is a Zariski closed subset of $\mathfrak{e}_{\Omega(n)} \times M_{\Omega}(n)$ and the maps

$$\pi_n : M_{\Omega}(n, \mathfrak{e}) \rightarrow M_{\Omega}(n) \quad \text{where } \pi_n(\nu, x) = x \quad (4.8)$$

and

$$\kappa_n : M_{\Omega}(n, \mathfrak{e}) \rightarrow \mathfrak{e}_{\Omega(n)} \quad \text{where } \kappa_n(\nu, x) = \nu \quad (4.9)$$

are surjective (see Theorem 2.5) algebraic morphisms.

For $m \in I_n$ and $i \in I_m$ let ρ_{im} be the regular function on $M_{\Omega}(n, \mathfrak{e})$ defined so that if $z \in M_{\Omega}(n, \mathfrak{e})$ and $\nu = \kappa_n(z)$ then $\rho_{im}(z) = \nu_{im}$.

Remark 4.1. One notes that if $z \in M_{\Omega}(n, \mathfrak{e})$ and $x \in M_{\Omega}(n)$ then $\pi_n(z) = x$ if and only if

$$(\rho_{1m}(z), \dots, \rho_{mm}(z)) = (\mu_{1m}(x), \dots, \mu_{mm}(x)) \quad (4.10)$$

up to a reordering, for all $m \in I_n$, using the notation of §2.2.

One defines a free action of Σ_n on $M_{\Omega}(n, \mathfrak{e})$, operating as a group of algebraic isomorphisms, by defining

$$\sigma \cdot z = (\sigma \cdot \nu, x) \quad (4.11)$$

where $\sigma \in \Sigma_n$ and $z = (\nu, x) \in M_\Omega(n, \mathfrak{e})$.

The following well-known proposition is classical.

Proposition 4.2. *Let $m \in I_n$ and let $\gamma(m) \in \mathfrak{e}(m)$. Let $g(m) \in Gl(m)$ and put $x(m) = Ad g(m)(\gamma(m))$. Then there exists an open neighborhood N of $\gamma(m)$ in $\mathfrak{e}(m)$ and a section $S \subset Gl(m)$ of the quotient map $Gl(m) \rightarrow Gl(m)/Diag(m)$ (using notation in §3.4) defined on a neighborhood of $g(m) Diag(m)$ such that the map*

$$S \times N \rightarrow M(m) \text{ where } (g, \gamma) \mapsto Ad g(\gamma) \quad (4.12)$$

is an analytic isomorphism onto an open set of $M(m)$. The elements of the image are necessarily regular semisimple elements of $M(m)$.

If U is an open subset of $M_n(\Omega)$ and $\phi : U \rightarrow \mathfrak{e}$ is an analytic map let $graph \phi : U \rightarrow \mathfrak{e} \times M_n(\Omega)$ be the analytic map defined by putting $graph \phi(x) = (\phi(x), x)$.

Proposition 4.3. *Let $z = (\nu, x) \in M_\Omega(n, \mathfrak{e})$. Then there exists a (sufficiently small) connected open neighborhood U of x in $M_\Omega(n)$ and an analytic map $\phi : U \rightarrow \mathfrak{e}_{\Omega(n)}$ with the following properties:*

(1) *$graph \phi(x) = z$, $graph \phi : U \rightarrow U_z$ is a homeomorphism, where U_z is the image of $graph \phi$, and $U_z \subset M_\Omega(n, \mathfrak{e})$.*

(2) *$\pi_n^{-1}(U) = \sqcup_{\sigma \in \Sigma_n} \sigma \cdot U_z$.*

(3) *$\{\sigma \cdot U_z \mid \sigma \in \Sigma_n\}$ are the connected components of $\pi_n^{-1}(U)$ and each component is open in $M_\Omega(n, \mathfrak{e})$.*

In particular U is evenly covered by π_n and π_n is a covering projection (see (4.8)).

Proof. Statements (1) and (2) are immediate consequences of Proposition 4.2. But it is immediate from (1) and (2) that $\sigma \cdot U_z$ is connected and closed in $\pi_n^{-1}(U)$ for any $\sigma \in \Sigma_n$. Since the partition in (2) is finite it follows that the parts are open in $\pi_n^{-1}(U)$ and hence are open in $M_\Omega(n, \mathfrak{e})$. The remaining statements are obvious. QED

4.3. As in the introduction, §0, let $\mathfrak{b}_e = -e + \mathfrak{b}$ using the notation of §2.2. By Remark 2.4, Theorems 2.3 and 2.5 hold if $\mathfrak{b}_e$ replaces $e + \mathfrak{b}$. Let $\mathfrak{b}_{e, \Omega(n)} = M_\Omega(n) \cap \mathfrak{b}_e$. Then $\mathfrak{b}_{e, \Omega(n)}$ is a Zariski open subset of $\mathfrak{b}_e$ and

$$\Phi_n : \mathfrak{b}_{e, \Omega(n)} \rightarrow \Omega(n) \quad (4.13)$$

is an algebraic isomorphism by Theorem 2.5 (see Remark 2.16). In particular $\mathfrak{b}_{e, \Omega(n)}$ is dense in $\mathfrak{b}_e$. Now let $M_\Omega(n, \mathfrak{e}, \mathfrak{b}) = \pi_n^{-1}(\mathfrak{b}_{e, \Omega(n)})$ so that, by Proposition 4.3, $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$

is a covering of $\mathfrak{b}_{e,\Omega(n)}$. Now consider the restriction

$$\kappa_n : M_\Omega(n, \mathfrak{e}, \mathfrak{b}) \rightarrow \mathfrak{e}_{\Omega(n)} \quad (4.14)$$

of (4.9) to $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$. Note that $\mathfrak{e}_{\Omega(n)}$ is connected since it is clearly Zariski open in $\mathfrak{e}$. One has

Theorem 4.4. *The map (4.14) is a homeomorphism. In particular $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ is connected.*

Proof. Let $\nu \in \mathfrak{e}_{\Omega(n)}$. Then by Theorem 2.5 there exists (uniquely) $x \in \mathfrak{b}_e$ such that, for any $m \in I_n$, $(\mu_{1m}(x), \dots, \mu_{m,m}(x)) = (\nu_{1,m}, \dots, \nu_{m,m})$, up to a reordering. But then $x \in \mathfrak{b}_{e,\Omega(n)}$ and, by Proposition 4.3, $(\nu, x) \in M_\Omega(n, \mathfrak{e}, \mathfrak{b})$. But then ν is in the image (4.14). That is, (4.14) is surjective. But assume $z, z' \in M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ and $\kappa_n(z) = \kappa_n(z')$. Then if $x = \pi_n(z)$ and $x' = \pi_n(z')$ one has $x, x' \in \mathfrak{b}_e$ and hence $x = x'$ by Theorem 2.5. Thus $z = z'$ so that (4.14) is injective. Hence (4.14) is bijective. But of course as a restriction map, (4.14) is continuous. We have only to show that its inverse is continuous.

Let $\beta_i, i \in I_n$, be the regular function on $\mathfrak{e}_{\Omega(n)}$ defined so that if $\nu \in \mathfrak{e}_{\Omega(n)}$ then $\beta_{d(m-1)+k}(\nu)$, $k \in I_m$, $m \in I_n$, is the elementary symmetric function of degree $m - k + 1$ in $\{\nu_{1m}, \dots, \nu_{mm}\}$. Now let $\beta : \mathfrak{e}_{\Omega(n)} \rightarrow \mathbb{C}^{d(n)}$ be the regular algebraic map defined so that

$$\beta(\nu) = (\beta_1(\nu), \dots, \beta_{d(n)}(\nu)) \quad (4.15)$$

One notes that if $c = \beta(\nu)$ then by (2.3),(2.4),(2.10) and (2.11)

$$(\nu_{1m}, \dots, \nu_{mm}) = (\mu_{1m}(c), \dots, \mu_{mm}(c)) \quad (4.16)$$

up to a reordering, for all $m \in I_n$. It follows then that

$$\beta : \mathfrak{e}_{\Omega(n)} \rightarrow \Omega(n) \quad (4.17)$$

is a surjective morphism (see (2.53)). Recalling Theorem 2.3 (where $-e$ replaces e), one has, inverting (4.13), a surjective morphism

$$\tilde{\beta} : \mathfrak{e}_{\Omega(n)} \rightarrow \mathfrak{b}_{e,\Omega(n)} \quad (4.18)$$

where $\beta = \Phi_n \circ \tilde{\beta}$ noting that, for $i \in I_n$,

$$p_i(\tilde{\beta}(\nu)) = \beta_i(\nu) \quad (4.19)$$

by (2.3),(2.4) and (2.5). But clearly $(\nu, \tilde{\beta}(\nu)) \in M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ for any $\nu \in \mathfrak{e}_{\Omega(n)}$. Hence

$$\mathfrak{e}_{\Omega(n)} \rightarrow M_\Omega(n, \mathfrak{e}, \mathfrak{b}), \quad \nu \mapsto (\nu, \tilde{\beta}(\nu)) \quad (4.20)$$

is an algebraic morphism. But (4.20) must be the inverse to (4.14), by the bijectivity of (4.14), since $\kappa_n((\nu, \tilde{\beta}(\nu))) = \nu$. Hence (4.14) is a homeomorphism. QED

Recalling (4.9) let $M_\nu(n, \mathfrak{e}) = \kappa_n^{-1}(\nu)$ so that one has a “fibration”

$$M_\Omega(n, \mathfrak{e}) = \sqcup_{\nu \in \mathfrak{e}_{\Omega(n)}} M_\nu(n, \mathfrak{e}) \quad (4.21)$$

of $M_\Omega(n, \mathfrak{e})$ over $\mathfrak{e}_{\Omega(n)}$ with fiber projection κ_n (see (4.9) and Proposition 4.5 below). Let $\nu \in \mathfrak{e}_{\Omega(n)}$. If $\nu \in \mathfrak{e}_{\Omega(n)}$ and $c \in \Omega(n) \subset \mathbb{C}^{d(n)}$ is defined by

$$c = \beta(\nu) \quad (4.22)$$

(see (4.17)) note that

$$M_c(n) \rightarrow M_\nu(n, \mathfrak{e}), \quad x \mapsto (\nu, x) \quad (4.23)$$

is a homeomorphism, by the definition of $M_\Omega(n, \mathfrak{e})$. The following asserts, in particular, that the “fibers” of (4.21) are all homeomorphic.

Proposition 4.5. *One has a homeomorphism*

$$M_\nu(n, \mathfrak{e}) \cong (\mathbb{C}^\times)^{d(n-1)} \quad (4.24)$$

for any $\nu \in \mathfrak{e}_{\Omega(n)}$.

Proof. This is immediate from (4.23) and Theorem 3.23. QED

Remark 4.6. Note that $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ defines a cross-section of κ_n by Theorem 4.14. That is, for any $\nu \in \mathfrak{e}_{\Omega(n)}$ the intersection

$$M_\Omega(n, \mathfrak{e}, \mathfrak{b}) \cap M_\nu(n, \mathfrak{e}) \text{ has only one point} \quad (4.25)$$

Proposition 4.7. *$M_\Omega(n, \mathfrak{e})$, as a topological space (Euclidean topology), is connected.*

Proof. As a covering of the manifold of $M_\Omega(n)$ obviously $M_\Omega(n, \mathfrak{e})$ is locally connected (see Proposition 4.3) so that any connected component of $M_\Omega(n, \mathfrak{e})$ is open in

$M_\Omega(n, \mathfrak{e})$. But then there exists a connected component C such that $M_\Omega(n, \mathfrak{e}, \mathfrak{b}) \subset C$, by Theorem 4.4. But if $\nu \in \mathfrak{e}_{\Omega(n)}$ then $M_\nu(n, \mathfrak{e})$ is connected by (4.24). But then $M_\nu(n, \mathfrak{e}) \subset C$ by (4.25). Hence $C = M_\Omega(n, \mathfrak{e})$ by (4.21). Thus $M_\Omega(n, \mathfrak{e})$ is connected. QED

4.4. The definition of variety here and throughout implies that it is Zariski irreducible. We will prove in this section that $M_\Omega(n, \mathfrak{e})$ is a nonsingular affine variety of dimension n^2 . We first observe

Proposition 4.8. *The nonempty Zariski open subset $\mathfrak{e}_{\Omega(n)}$ of $\mathfrak{e}$ (see §4.1) is a nonsingular affine variety and the nonempty Zariski open subset $M_\Omega(n)$ of $M(n)$ is a nonsingular affine variety (see (2.53)). In particular $\mathfrak{e}_{\Omega(n)} \times M_\Omega(n)$ is a nonsingular affine variety of dimension $n^2 + d(n)$.*

Proof. Recalling §4.1, clearly, for any $m \in I_n$, $\mathfrak{e}(m)$ is a Zariski open, nonempty subvariety of $\mathfrak{d}(m)$. It is affine since it is the complement of the zero set of the discriminant function on $\mathfrak{d}(m)$. But then $\mathfrak{e}$ is a nonsingular affine variety of dimension $d(n)$. But then $\mathfrak{e}_{\Omega(n)}$ is a nonsingular affine variety of dimension $d(n)$ since the condition (4.4) clearly defines $\mathfrak{e}_{\Omega(n)}$ as the complement of the zero set of a single regular function on $\mathfrak{e}$. But now the argument in Remark 2.16 readily characterizes $M_\Omega(n)$ as the complement in $M(n)$ of the zero set in $M(n)$ of a polynomial in $J(n)$ (see (2.30)). Thus $M_\Omega(n)$ is a nonsingular affine variety of dimension n^2 . This of course proves the proposition. QED

If X is an affine variety we will denote the affine ring of X by $\mathcal{O}(X)$.

Theorem 4.9. *$M_\Omega(n, \mathfrak{e})$ is a n^2 -dimensional Zariski closed affine nonsingular subvariety of the nonsingular affine variety $\mathfrak{e}_{\Omega(n)} \times M_\Omega(n)$ (see Proposition 4.8).*

Proof. One has

$$\mathcal{O}(\mathfrak{e}_{\Omega(n)} \times M_\Omega(n)) = \mathcal{O}(\mathfrak{e}_{\Omega(n)}) \otimes \mathcal{O}(M_\Omega(n)) \quad (4.26)$$

For $i \in I_n$ let $\beta_i \in \mathcal{O}(\mathfrak{e}_{\Omega(n)})$ be defined as in (4.15). Also let $p'_i \in \mathcal{O}(M_\Omega(n))$ be defined by putting $p'_i = p_i|_{M_\Omega(n)}$ where $p_i \in J(n)$ is given by (2.5) (see (2.30)). For notational convenience put $W = \mathfrak{e}_{\Omega(n)} \times M_\Omega(n)$. Now let $f_i \in \mathcal{O}(W)$ be defined by putting $f_i = \beta_i \otimes 1 - 1 \otimes p'_i$. Clearly

$$M_\Omega(n, \mathfrak{e}) = \text{Spec}[\mathcal{O}(W)/(f_1, \dots, f_{d(n)})] \quad (4.27)$$

But for any $z = (\nu, x) \in M_\Omega(n, \mathfrak{e})$ one has

$$(df_i)_z, i \in I_n, \text{ are linearly independent} \quad (4.28)$$

since $(dp_i)_x, i \in I_n$, are linearly independent in $T_x^*(M(n))$ by (2.55) and the definition of $M^{sreg}(n)$ in §2.3. One also notes that $(d\beta_i)_\nu, i \in I_n$, are linearly independent in $T_\nu^*(\mathfrak{e})$ since $\nu(m)$ is regular in $\mathfrak{d}(m)$ for any $m \in I_n$. But then $M_\Omega(n, \mathfrak{e})$ is nonsingular at z by Theorem 4 in §4 of Chapter 3 in [M1], p. 172, where X and U in the notation of that reference are equal to W here (see Proposition 4.8) and $Y = M_\Omega(n, \mathfrak{e})$. Thus $M_\Omega(n, \mathfrak{e})$ is nonsingular and has dimension n^2 . But now since $M_\Omega(n, \mathfrak{e})$ is connected in the Euclidean topology, by Proposition 4.7, it is obviously connected in the Zariski topology (i.e. it is not the disjoint union of two nonempty Zariski open sets). But then, since it is nonsingular, it is irreducible as an algebraic set by Corollary 17.2, p. 72, in [B]. It is then also a closed affine subvariety of W by (4.27). QED

We may regard $\mathcal{O}(M_\Omega(n))$ as a module for $J(n)$ (see §2.4)) where p_i operates as multiplication by p'_i (using the notation in the proof of Theorem 4.9). Similarly regard $\mathcal{O}(\mathfrak{e}_{\Omega(n)})$ as a module for $J(n)$ where p_i operates as multiplication by β_i (see (4.19)). Then Theorem 4.9 and the equality (4.27) clearly implies

Theorem 4.10. *As an affine ring one has the tensor product*

$$\mathcal{O}(M_\Omega(n, \mathfrak{e})) = \mathcal{O}(\mathfrak{e}_{\Omega(n)}) \otimes_{J(n)} \mathcal{O}(M_\Omega(n)) \quad (4.29)$$

4.5. Assume Y is an affine variety and that Σ is a finite group operating as a group of algebraic isomorphisms of Y . Then Σ operates as a group of algebraic automorphisms of $\mathcal{O}(Y)$ where for any $f \in \mathcal{O}(Y)$, $\sigma \in \Sigma$ and $z \in Y$, one has

$$(\sigma \cdot f)(z) = f(\sigma^{-1} \cdot z) \quad (4.30)$$

Let Y/Σ be the set of orbits of Σ on Y . Obviously any $f \in \mathcal{O}(Y)^\Sigma$ defines a function on Y/Σ . It is then a classical theorem that Y/Σ has the structure of an affine variety where

$$\mathcal{O}(Y/\Sigma) = \mathcal{O}(Y)^\Sigma \quad (4.31)$$

See e.g. Theorem 1.1, p. 27 and Amplification 1.3, p. 30 in Chapter 1, §2 of [M2]. In addition one notes that

$$Y \rightarrow Y/\Sigma, z \mapsto \Sigma \cdot z \quad (4.32)$$

is a morphism where the corresponding cohomorphism is the embedding

$$\mathcal{O}(Y)^\Sigma \rightarrow \mathcal{O}(Y) \quad (4.33)$$

Let $F(Y)$ be the quotient field of $\mathcal{O}(Y)$ and let $F(Y/\Sigma)$ be the quotient field of $\mathcal{O}(Y/\Sigma)$. Since Σ is finite it is a simple and well-known fact that, as a consequence of (4.31),

$$F(Y/\Sigma) = F(Y)^\Sigma \text{ so that } F(Y) \text{ is a Galois extension of } F(Y/\Sigma) \quad (4.34)$$

Let $Clos_{F(Y)}(\mathcal{O}(Y/\Sigma))$ be the integral closure of $\mathcal{O}(Y/\Sigma)$ in $F(Y)$.

Proposition 4.11. *$Clos_{F(Y)}(\mathcal{O}(Y/\Sigma))$ is a finite module over $\mathcal{O}(Y/\Sigma)$. In particular $Clos_{F(Y)}(\mathcal{O}(Y/\Sigma))$ is Noetherian. Furthermore $\mathcal{O}(Y)$ is also a finite module over $\mathcal{O}(Y/\Sigma)$ so that $\mathcal{O}(Y)$ is integral over $\mathcal{O}(Y/\Sigma)$ and hence*

$$\mathcal{O}(Y) \subset Clos_{F(Y)}(\mathcal{O}(Y/\Sigma)) \quad (4.35)$$

and one has equality in (4.35) in case Y is nonsingular. Finally, (in the sense of Chapter 2, §7, Definition 3, p. 124 in [M1]) the morphism (4.32) is finite and the morphism

$$Z \mapsto Y/\Sigma \quad (4.36)$$

is finite where $Z = \text{Spec}(Clos_{F(Y)}(\mathcal{O}(Y/\Sigma)))$ and (4.36) is defined so that the injection $\mathcal{O}(Y/\Sigma) \rightarrow Clos_{F(Y)}(\mathcal{O}(Y/\Sigma))$ is the corresponding cohomomorphism.

Proof. The first statement is given by Theorem 9, in Chapter 5, §4, p.267 in [ZS]. The statement that $\mathcal{O}(Y)$ is also a finite module over $\mathcal{O}(Y/\Sigma)$ is stated as Noether's Theorem and proved as Theorem 2.3.1, p. 26 in [Sm]. But now if Y is nonsingular then $\mathcal{O}(Y)$ is integrally closed in $F(Y)$. See e.g. the first paragraph, p. 197, in [M1]. But of course $Clos_{F(Y)}(\mathcal{O}(Y/\Sigma))$ is integral over $\mathcal{O}(Y)$. Hence one has equality in (4.35). But now the finiteness of (4.32) and (4.36) follows from Proposition 5, p. 124 in §7 of Chapter 2 in [M1] since $Y, Y/\Sigma$ and Z are affine varieties. QED

Making use of Theorem 4.9 we apply Proposition 4.11 in the case where $Y = M_\Omega(n, \mathfrak{e})$ and $\Sigma = \Sigma_n$ with, of course, the action given by (4.12). The quotient field of $\mathcal{O}(M_\Omega(n, \mathfrak{e}))$ will be denoted by $F(n, \mathfrak{e})$. The quotient field of $\mathcal{O}(M(n))$ will be denoted by $F(n)$. In the notation of §1.1 note that $\mathcal{O}(M(n))$ is just $P(n)$. Since $M_\Omega(n)$ is Zariski dense in $M(n)$ note that $F(n)$ is also the quotient field of $\mathcal{O}(M_\Omega(n))$.

It is clear, from the definition of π_n (see (4.8)), that π_n is a morphism whose corresponding cohomomorphism maps $\mathcal{O}(M_\Omega(n))$ into $\mathcal{O}(M_\Omega(n, \mathfrak{e})^{\Sigma_n})$. Hence π_n descends to a morphism

$$M_\Omega(n, \mathfrak{e})^{\Sigma_n} \rightarrow M_\Omega(n) \quad (4.37)$$

Proposition 4.12. *The morphism (4.37) is an isomorphism of algebraic varieties. That is,*

$$M_{\Omega}(n, \mathfrak{e})^{\Sigma_n} \cong M_{\Omega}(n) \quad (4.38)$$

In particular π_n is a finite morphism. Furthermore

$$F(n, \mathfrak{e})^{\Sigma_n} \cong F(n) \quad (4.39)$$

so that, using (4.39) to define an identification, $F(n, \mathfrak{e})$ is a Galois extension of $F(n)$ with Galois group Σ_n . In addition, using (4.38) to define an identification one has

$$\text{Clos}_{F(n, \mathfrak{e})} \mathcal{O}(M_{\Omega}(n)) = \mathcal{O}(M_{\Omega}(n, \mathfrak{e})) \quad (4.40)$$

Proof. It is immediate from Proposition 4.3 that (4.37) is bijective. But then it is birational by Theorem 5.1.6, p. 81 in [Sp] (since we are in a characteristic zero case). But $M_{\Omega}(n)$ is nonsingular. Thus (4.37) is an isomorphism (see e.g. Theorem 5.2.8, p. 85 in [Sp]). The rest of the statements follow from Proposition 4.11 since $M_{\Omega}(n, \mathfrak{e})$ is nonsingular by Theorem 4.9. QED

Recall the notation of Proposition 4.3 so that $z = (\gamma, x) \in M_{\Omega}(n, \mathfrak{e})$. Also U_z is a (Euclidean) open neighborhood of z in $M_{\Omega}(n, \mathfrak{e})$, U is a (Euclidean) open neighborhood of x in $M_{\Omega}(n)$ and the statements of Proposition 4.3 hold. The inverse of the homeomorphism *graph* $\phi : U \rightarrow U_z$ is clearly

$$\pi_n|_{U_z} : U_z \rightarrow U \quad (4.41)$$

Proposition 4.13. *Recalling the notation of Proposition 4.3 the map (4.41) is an analytic isomorphism. In particular (see Corollary 2, p. 182, to Theorem 3 in §5 of Chapter 3 in [M1]) $\pi_n : M_{\Omega}(n, \mathfrak{e}) \rightarrow M_{\Omega}(n)$ (see (4.8)) is an étale morphism.*

Proof. Since π_n is a morphism it is a holomorphic map of nonsingular analytic manifolds (see ii, p. 58 of §10, Chapter 1 in [M1]) Thus the homeomorphism (4.41) is an analytic map. It suffices to prove that

$$\text{graph } \phi : U \rightarrow U_z \quad (4.42)$$

is analytic. To do this first regard (4.42) as a map

$$U \rightarrow \mathfrak{e}_{\Omega(n)} \times M_{\Omega}(n) \quad (4.43)$$

(see (4.7)). By the definition of $\text{graph } \phi$ in Proposition 4.3 it is obvious that (4.43) is analytic. Hence if $g \in \mathcal{O}(\mathfrak{e}_{\Omega(n)} \times M_{\Omega}(n))$ then $g \circ \text{graph } \phi$ is analytic function on U . But then $f \circ \text{graph } \phi$ is an analytic function on U for any $f \in \mathcal{O}(M_{\Omega}(n, \mathfrak{e}))$ since $\mathcal{O}(M_{\Omega}(n, \mathfrak{e}))$ is just the restriction of $\mathcal{O}(\mathfrak{e}_{\Omega(n)} \times M_{\Omega}(n))$ to $M_{\Omega}(n, \mathfrak{e})$. But then (4.42) is analytic since an analytic coordinate system in a Euclidean neighborhood of z in $M_{\Omega}(n, \mathfrak{e})$ is given by elements in $\mathcal{O}(M_{\Omega}(n, \mathfrak{e}))$ which are uniformizing parameters in a Zariski neighborhood of z (see p. 183, §6 of Chapter 3 in [M1]). QED

Combining Propositions 4.3 and 4.13 one has

Proposition 4.14. *The map π_n defines $M_{\Omega}(n, \mathfrak{e})$ an analytic covering of $M_{\Omega}(n)$ with Σ_n as the group of deck transformations.*

4.6. Since $M_{\Omega}(n, \mathfrak{e})$ is locally and analytically isomorphic to $M_{\Omega}(n)$ (via π_n) the tensor which defines Poisson bracket of functions on $M_{\Omega}(n)$ lifts and defines Poisson bracket of analytic functions on $M_{\Omega}(n, \mathfrak{e})$. In particular $\mathcal{O}(M_{\Omega}(n, \mathfrak{e}))$ has the structure of a Poisson algebra. For any $f \in \mathcal{O}(M_{\Omega}(n))$ (noting that we regard $\mathcal{O}(M(n)) \subset \mathcal{O}(M_{\Omega}(n))$) let $\widehat{f} \in \mathcal{O}(M_{\Omega}(n, \mathfrak{e}))$ be defined by putting $\widehat{f} = f \circ \pi_n$. For $f_1, f_2 \in \mathcal{O}(M_{\Omega}(n))$ one then has

$$[\widehat{f_1}, \widehat{f_2}] = \widehat{[f_1, f_2]} \quad (4.44)$$

Using the notation of §1.2 but now, in addition, applied to $M_{\Omega}(n, \mathfrak{e})$, for any $\varphi \in \mathcal{O}(M_{\Omega}(n, \mathfrak{e}))$ let ξ_{φ} be the (complex) vector field on $M_{\Omega}(n, \mathfrak{e})$ defined so that $\xi_{\varphi}\psi = [\varphi, \psi]$ for any $\psi \in \mathcal{O}(M_{\Omega}(n))$. It is immediate that if $f \in \mathcal{O}(M_{\Omega}(n))$ then $\xi_{\widehat{f}}$ is π_n -related to ξ_f so that unambiguously

$$(\pi_n)_*(\xi_{\widehat{f}}) = \xi_f \quad (4.45)$$

Besides the (just considered) subring of $\mathcal{O}(M_{\Omega}(n, \mathfrak{e}))$, defined by the pull-back of (the surjection) π_n , there is the subring of $\mathcal{O}(M_{\Omega}(n, \mathfrak{e}))$ defined by the pull-back of (the surjection) κ_n (see (4.9)). Indeed let

$$J(n, \mathfrak{e}) = \{q \circ \kappa_n \mid q \in \mathcal{O}(\mathfrak{e}_{\Omega(n)})\} \quad (4.46)$$

From the definition of ρ_{km} , $k \in I_m$, $m \in I_m$ in §4.2 note that $\rho_{km} \in J(n, \mathfrak{e})$. For notational convenience let $r_i \in J(n, \mathfrak{e})$, $i \in I_{d(n)}$ be defined so that

$$r_i = \rho_{km} \quad (4.47)$$

where

$$i = d(m-1) + k \quad (4.48)$$

so that

$$r_i \in J(n, \mathfrak{e}), \quad i \in d(n) \quad (4.49)$$

Remark 4.15. Recalling §4.1 note that conversely $J(n, \mathfrak{e})$ is a localization of the polynomial ring generated by the r_i , $i \in I_{d(n)}$.

Now put $\widehat{J(n)} = \{\widehat{p} \mid p \in J(n)\}$ (see §2.4) so that $\widehat{J(n)}$ is the polynomial ring

$$\widehat{J(n)} = \mathbb{C}[\widehat{p}_1, \dots, \widehat{p}_{d(n)}] \quad (4.50)$$

Let $I_{[m]} = I_{d(m)} - I_{d(m-1)}$ so that $\text{card } I_{[m]} = m$.

Proposition 4.16. *One has*

$$\widehat{J(n)} \subset J(n, \mathfrak{e}) \quad (4.51)$$

In fact if $i \in I_{[m]}$, where $m \in I_n$, and i is written as in (4.48) then $\widehat{p}_i$ is the elementary symmetric polynomial of degree $m - k + 1$ in the functions r_j , $j \in I_{[m]}$. Indeed if

$$P_m(\lambda) = \lambda^m + \sum_{k \in I_m} (-1)^{m-k+1} p_{d(m-1)+k} \lambda^{k-1}$$

then

$$P_m(\lambda) = \prod_{j \in I_{[m]}} (\lambda - r_j) \quad (4.52)$$

so that, in addition, r_j , for $j \in I_{[m]}$, satisfies the polynomial equation

$$P_m(r_j) = 0$$

Proof. The inclusion (4.51) follows from (4.50) and (4.52). On the other hand (4.52) follows from (2.3), (2.4) and (2.5) together with (4.10), (4.29) and (4.47). QED

Since π_n is an analytic covering map (see Proposition 4.14) it follows from (2.55) and the definition of strongly regular (see §2.3) that the differentials $(d\widehat{p}_i)_z$, $i \in I_{d(n)}$, are linearly independent at any $z \in M_\Omega(n, \mathfrak{e})$. For any $m \in I_n$ let $T_z^*(M_\Omega(n, \mathfrak{e}))^{(m)}$ be the m -dimensional subspace of the cotangent space $T_z^*(M_\Omega(n, \mathfrak{e}))$ spanned by the differentials $(d\widehat{p}_i)_z$, $i \in I_{[m]}$.

Proposition 4.17. *Let $z \in M_\Omega(n, \mathfrak{e})$ and let $m \in I_n$. Then $(dr_i)_z, i \in I_{[m]}$ is a basis of $T_z^*(M_\Omega(n, \mathfrak{e}))^{(m)}$.*

Proof. Let V be the space of $T_z^*(M_\Omega(n, \mathfrak{e}))$ spanned by $(dr_i)_z, i \in I_{[m]}$. But then $\dim V \leq m$. But $T_z^*(M_\Omega(n, \mathfrak{e}))^{(m)} \subset V$ by (4.51). Thus $V = T_z^*(M_\Omega(n, \mathfrak{e}))^{(m)}$ by dimension. QED

In the notation above let $T_z^*(M_\Omega(n, \mathfrak{e}))'$ (resp. $T_z^*(M_\Omega(n, \mathfrak{e}))''$) be the $d(n-1)$ - (resp $d(n)$ -) dimensional sum of the subspaces $T_z^*(M_\Omega(n, \mathfrak{e}))^{(m)}$ over all $m \in I_{n-1}$ (resp. $m \in I_n$). Then as an immediate consequence of Proposition 4.17 one has

Proposition 4.18. *Let $z \in M_\Omega(n, \mathfrak{e})$. Then $(dr_i)_z, i \in I_{d(n-1)}$ (resp. $I_{d(n)}$) is a basis of $T_z^*(M_\Omega(n, \mathfrak{e}))'$ (resp. $T_z^*(M_\Omega(n, \mathfrak{e}))''$).*

4.7. Let $\nu \in \mathfrak{e}_{\Omega(n)}$. By definition (see (4.21)) $M_\nu(n, \mathfrak{e}) = \kappa_n^{-1}(\nu)$ so that $M_\nu(n, \mathfrak{e})$ is a Zariski closed subset of $M_\Omega(n, \mathfrak{e})$.

Remark 4.19. Note that (see (4.2), (4.3), §4.2 and (4.47)) $M_\nu(n, \mathfrak{e})$ may be given by the equations

$$M_\nu(n, \mathfrak{e}) = \{z \in M_\Omega(n, \mathfrak{e}) \mid r_i(z) = \nu_{k m} \text{ when } i \in I_{d(n)} \text{ is put in the form (4.48)}\} \quad (4.53)$$

Note also that $M_\nu(n, \mathfrak{e})$ is nonsingular by Proposition 4.18 (linear independence of differentials).

Proposition 4.20. *Let $\nu \in \mathfrak{e}_{\Omega(n)}$ and let $c = \beta(\nu)$ so that $c \in \Omega(n)$ (see (4.22)). Then the covering map π_n (see (4.8)) restricts to an algebraic isomorphism*

$$\pi_n : M_\nu(n, \mathfrak{e}) \rightarrow M_c(n) \quad (4.54)$$

of nonsingular affine varieties.

Proof. Recall that $M_c(n)$ is an irreducible nonsingular Zariski closed subvariety of $M(n)$ (see Theorem 3.23). The homeomorphism (4.23) can obviously be regarded as a morphism mapping $M_c(n)$ to $\mathfrak{e}_{\Omega(n)} \times M_\Omega(n)$. However the image of (4.23) is the Zariski closed subset $M_\nu(n, \mathfrak{e})$ of $\mathfrak{e}_{\Omega(n)} \times M_\Omega(n)$. Thus (4.23), as it stands, is a bijective morphism. But then $M_\nu(n, \mathfrak{e})$ is a variety (i.e. it is irreducible). Hence it is a nonsingular affine variety by Remark 4.19. Thus (4.23), as it stands, is an algebraic isomorphism. But (4.54) is just the inverse of (4.23). QED

Note that (2.31) and (4.44) imply

$$[\widehat{p}, \widehat{q}] = 0 \quad (4.55)$$

for any $p, q \in J(n)$. In particular

$$[\widehat{p}_i, \widehat{p}_j] = 0 \quad (4.56)$$

for any $i, j \in I_{d(n)}$. One consequence of Proposition 4.19 is

Proposition 4.21. *Let $\nu \in \mathfrak{e}_{\Omega(n)}$ and let $z \in M_\nu(n, \mathfrak{e})$, so that $\nu = \kappa_n(z)$ (see (4.9)). Then $(\xi_{\widehat{p}_i})_z$, $i \in I_{d(n-1)}$, is a basis of the tangent space $T_z(M_\nu(n, \mathfrak{e}))$.*

Proof. Let $x = \pi_n(z)$ (see (4.8)) and $c = \beta(\nu)$ (see (4.22)) so that $x \in M_c(n)$. Since π_n is a local analytic isomorphism it suffices by Proposition 4.20 and (4.45) to see that $(\xi_{p_i})_x$, $i \in d(n-1)$, is a basis of $T_x(M_c(n))$. But x is strongly regular (see §2.3) by (2.55) since $c \in \Omega(n)$. The result then follows from Remark 2.8 and Theorems 3.4, 3.23. QED

The argument which established Theorem 3.25 may now be used to establish

Theorem 4.22. *$J(n, \mathfrak{e})$ (see (4.46)) is a maximal Poisson commutative subalgebra of $\mathcal{O}(M_\Omega(n, \mathfrak{e}))$. In particular (see (4.48))*

$$[r_i, r_j] = 0 \quad (4.57)$$

for any $i, j \in I_{d(n)}$. Furthermore if $f \in J(n, \mathfrak{e})$ and $\nu \in \mathfrak{e}_{\Omega(n)}$ then $f|_{M_\nu(n, \mathfrak{e})}$ is a constant function and $\xi_f|_{M_\nu(n, \mathfrak{e})}$ is tangent to $M_\nu(n, \mathfrak{e})$. Moreover if we write (using Proposition 4.21)

$$\xi_f = \sum_{i \in I_{d(n-1)}} f_i \xi_{\widehat{p}_i} \quad (4.58)$$

on $M_\nu(n, \mathfrak{e})$ where $f_i \in \mathcal{O}(M_\nu(n, \mathfrak{e}))$ then all the f_i are constant on $M_\nu(n, \mathfrak{e})$. Finally

$$\begin{aligned} (\xi_{r_i})_z, i \in I_{d(n-1)}, \text{ is a basis of } T_z(M_\nu(n, \mathfrak{e})) \text{ for any } z \in M_\nu(n, \mathfrak{e}) \text{ and} \\ (\xi_{r_i})_z = 0, i \in I_{[n]} = I_{d(n)} - I_{d(n-1)}, \text{ for any } z \in M_\nu(n, \mathfrak{e}) \end{aligned} \quad (4.59)$$

Proof. Let $\nu \in \mathfrak{e}_{\Omega(n)}$. The function r_i is constant on $M_\nu(n, \mathfrak{e})$ for all $i \in I_{d(n)}$ by Remark 4.19. Let $f \in J(n, \mathfrak{e})$. But then $f|_{M_\nu(n, \mathfrak{e})}$ is a constant function by Remark 4.15. On the other hand if $z \in M_\nu(n, \mathfrak{e})$ and W_z is the orthocomplement of

$T_z(M_\nu(n, \mathfrak{e}))$ in $T^*(M_\Omega(n, \mathfrak{e}))$ then $(dr_i)_z$, $i \in I_{d(n)}$, is a basis of W_z by Proposition 4.18 and Remark 4.19. But this implies that $(df)_z \in W_z$ by Remark 4.15. In particular $(d\widehat{p}_i)_z \in W_z$ for any $i \in I_{d(n)}$ by (4.50). In fact $(d\widehat{p}_i)_z$, $i \in I_{d(n)}$, is a basis of W_z by Proposition 4.18 and the definition of $T_z^*(M_\Omega(n, \mathfrak{e}))''$ in §4.6.

Now for any $g \in \mathcal{O}(M_\Omega(n, \mathfrak{e}))$ the tangent vector $(\xi_g)_z$ depends only on $(dg)_z$ (see §1.2). But

$$(\xi_{\widehat{p}_i})_z = 0 \text{ for any } i \in I_{[n]} = I_{d(n)} - I_{d(n-1)} \quad (4.60)$$

by (2.7). But then $(\xi_g)_z \in T_z(M_\nu(n, \mathfrak{e}))$ if $(dg)_z \in W_z$, by Proposition 4.21. Hence $\xi_f|_{M_\nu(n, \mathfrak{e})}$ is tangent to $M_\nu(n, \mathfrak{e})$. Furthermore Propositions 4.17, 4.18 and 4.21 imply (4.59). Now if $g \in J(n, \mathfrak{e})$ then $g|_{M_\nu(n, \mathfrak{e})}$ is constant. But $\xi_f|_{M_\nu(n, \mathfrak{e})}$ is tangent to $M_\nu(n, \mathfrak{e})$. Thus $J(n, \mathfrak{e})$ is Poisson commutative. In particular $[\xi_{\widehat{p}_j}, \xi_f] = 0$ for any $j \in I_{d(n-1)}$. But, on $M_\nu(n, \mathfrak{e})$, one has

$$[\xi_{\widehat{p}_j}, \xi_f] = \sum_{i \in I_{d(n-1)}} (\xi_{\widehat{p}_j} f_i) \xi_{\widehat{p}_i}$$

by (4.58). Thus $\xi_{\widehat{p}_j} f_i = 0$, by Proposition 4.21, for all $i, j \in I_{d(n-1)}$. Hence the f_i are constants.

Now recall the definition of $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ in §4.3. Then $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ is a Zariski closed subset of $M_\Omega(n, \mathfrak{e})$ since, clearly, $\mathfrak{b}_{\mathfrak{e}, \Omega(n)}$ is obviously closed in $M_\Omega(n)$. But $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ is irreducible since it is the image of the bijective morphism (4.20). But then (4.14) is a bijective (and hence, necessarily birational, since we are in characteristic 0) morphism of irreducible varieties. In addition $\mathfrak{e}_{\Omega(n)}$ is nonsingular (see §4.1). Thus (4.14) is an isomorphism of varieties. Consequently (see (4.46)) the map

$$J(n, \mathfrak{e}) \rightarrow \mathcal{O}(M_\Omega(n, \mathfrak{e}, \mathfrak{b})), \quad g \mapsto g|_{M_\Omega(n, \mathfrak{e}, \mathfrak{b})} \quad (4.61)$$

is an algebra isomorphism. Consequently given any $h \in \mathcal{O}(M_\Omega(n, \mathfrak{e}))$ there exists a unique $g \in J(n, \mathfrak{e})$ such that $g|_{M_\Omega(n, \mathfrak{e}, \mathfrak{b})} = h|_{M_\Omega(n, \mathfrak{e}, \mathfrak{b})}$. But now assume that h poisson commutes with any function in $J(n, \mathfrak{e})$. Then $h|_{M_\nu(n, \mathfrak{e})}$ is constant, by Proposition 4.21, for any $\nu \in \mathfrak{e}_{\Omega(n)}$. But then $h = g$ on $M_\nu(n, \mathfrak{e})$ by (4.25). Thus $h = g$, by (4.21), and hence $J(n, \mathfrak{e})$ is maximally Poisson commutative in $\mathcal{O}(M_\Omega(n, \mathfrak{e}))$. QED

4.8. We recall some definitions, results and notations in Part 1. Generators $p_{(i)}$, $i \in I_{d(n)}$, of the polynomial ring $J(n)$ (see (2.30)) were defined by (3.20), recalling (2.38). In particular two sets of generators of $J(n)$ were under consideration in Part 1; namely the $p_{(i)}$ and the p_i (see (2.5) and (2.3)) where $i \in I_{d(n)}$. From the discussion preceding (2.38) it follows that for $x \in M(n)$, and $m \in I_n$

$$\text{span of } (dp_{(i)})_x, \text{ for } i \in I_{d(m)} = \text{span of } (dp_i)_x, \text{ for } i \in I_{d(m)} \quad (4.62)$$

and hence

$$\text{span of } (\xi_{p(i)})_x, \text{ for } i \in I_{d(m)} = \text{span of } (\xi_{p_i})_x, \text{ for } i \in I_{d(m)} \quad (4.63)$$

But then, by (4.45), for any $z \in M_\Omega(n, \mathfrak{e})$,

$$\text{span of } (\xi_{p(i)}^\frown)_z, \text{ for } i \in I_{d(m)} = \text{span of } (\xi_{p_i}^\frown)_z, \text{ for } i \in I_{d(m)} \quad (4.64)$$

An immediate consequence of (4.64), when $m = n - 1$, and Proposition 4.21 is

Proposition 4.23. *Let $\nu \in \mathfrak{e}_{\Omega(n)}$ and let $z \in M_\nu(n, \mathfrak{e})$. Then $(\xi_{p(i)}^\frown)_z, i \in I_{d(n-1)}$, is a basis of $T_z(M_\nu(n, \mathfrak{e}))$.*

By definition $\mathfrak{a}$ (see §3.2) is the (complex) commutative $d(n - 1)$ -dimensional Lie algebra of vector fields on $M(n)$ spanned by $\xi_{p(i)}, i \in I_{d(n-1)}$. By Theorem 3.4 the Lie algebra $\mathfrak{a}$ integrates to a (complex) analytic Lie group $A \cong \mathbb{C}^{d(n-1)}$ which operates analytically on $M(n)$. Now let $\hat{\mathfrak{a}}$ be the $d(n - 1)$ -dimensional complex commutative (see (4.44)) Lie algebra of vector fields on $M_\Omega(n, \mathfrak{e})$ spanned by $\xi_{p(i)}^\frown, i \in I_{d(n-1)}$.

Remark 4.24. If $\nu \in \mathfrak{e}_{\Omega(n)}$ then note that, by Proposition 4.23, $\hat{\mathfrak{a}}|_{M_\nu(n, \mathfrak{e})}$ is a commutative $d(n - 1)$ -dimensional Lie algebra of vector fields on $M_\nu(n, \mathfrak{e})$.

Let $\hat{A} (\cong \mathbb{C}^{d(n-1)})$ be a simply connected Lie group with Lie algebra $\hat{\mathfrak{a}}$. Let

$$A \rightarrow \hat{A}, a \mapsto \hat{a} \quad (4.65)$$

be the group isomorphism whose differential maps $\xi_{p(i)}$ to $\xi_{p(i)}^\frown$ for all $i \in I_{d(n-1)}$.

Theorem 4.25. *The Lie algebra $\hat{\mathfrak{a}}$ integrates to an action of $\hat{A}$ on $M_\Omega(n, \mathfrak{e})$. Furthermore if $a \in A$ and $z \in M_\Omega(n, \mathfrak{e})$ then*

$$\pi_n(\hat{a} \cdot z) = a \cdot x \quad (4.66)$$

where $x = \pi_n(z)$. Moreover $M_\nu(n, \mathfrak{e})$ is stable under the action of $\hat{A}$ for any $\nu \in \mathfrak{e}_{\Omega(n)}$. In fact, for all ν , $\hat{A}$ operates transitively on $M_\nu(n, \mathfrak{e})$ so that (4.21) is the decomposition of $M_\Omega(n, \mathfrak{e})$ into $\hat{A}$ orbits.

Proof. Noting Remark 4.24, Theorem 4.24 is an immediate consequence of (4.45), the isomorphism (4.54) and Theorem 3.23. QED

Remark 4.26. Implicit in Theorem 4.25 and its proof is the fact that if $\nu \in \mathfrak{e}_{\Omega(n)}$ then the Lie algebra, $\widehat{\mathfrak{a}}|M_\nu(n, \mathfrak{e})$ of vector fields on $M_\nu(n, \mathfrak{e})$ (see Remark 4.24) integrates to the group action $\widehat{A}|M_\nu(n, \mathfrak{e})$ on $M_\nu(n, \mathfrak{e})$.

One now has an analogue of Theorem 3.5. (Actually it is analogue of a considerably weaker result than Theorem 3.5 in that $M_\Omega(n, \mathfrak{e})$ covers the strongly regular set $M_\Omega(n)$ and not all of $M(n)$.)

Theorem 4.27. *Let $f \in J(n, \mathfrak{e})$ (see (4.46)). Then the vector field ξ_f integrates to an action of $\mathbb{C}$ on $M_\Omega(n, \mathfrak{e})$. In fact if $\nu \in \mathfrak{e}_{\Omega(n)}$ then $\xi_f|_{M_\nu(n, \mathfrak{e})}$ is tangent to $M_\nu(n, \mathfrak{e})$. Indeed*

$$\xi_f|_{M_\nu(n, \mathfrak{e})} \in \widehat{\mathfrak{a}}|_{M_\nu(n, \mathfrak{e})} \quad (4.67)$$

so that (see Remark 4.26) the action of $\mathbb{C}$ stabilizes $M_\nu(n, \mathfrak{e})$.

Proof. Clearly $\widehat{p_{(i)}} \in J(n, \mathfrak{e})$, for $i \in I_{d(n-1)}$, by (4.50). Let $\nu \in \mathfrak{e}_{\Omega(n)}$. Then $\xi_{\widehat{p_{(i)}}}|_{M_\nu(n, \mathfrak{e})}$, $i \in I_{d(n-1)}$, is a basis of $\widehat{\mathfrak{a}}|_{M_\nu(n, \mathfrak{e})}$ by (the constancy of the f_i in) Theorem 4.22 and Proposition 4.23. But then one has (4.67), also by Theorem 4.22. Theorem 4.27 then follows from Remark 4.26 and Theorem 4.25. QED

5. The Emergence of the dual coordinates s_j , $j \in I_{d(n-1)}$

5.1. We first wish to be more explicit about the vector fields ξ_{r_j} , $j \in I_{d(n-1)}$ on $M_\Omega(n, \mathfrak{e})$. See §4.2 and (4.47). Fix $m \in I_n$. We have put $I_{[m]} = I_{d(m)} - I_{d(m-1)}$. For $i \in I_{[m]}$ and

$$i = d(m-1) + k \quad (5.1)$$

for $k \in I_m$ one has, on $M_\Omega(n, \mathfrak{e})$,

$$\widehat{p_{(i)}} = \frac{1}{m+1-k} \sum_{j \in I_{[m]}} r_j^{m+1-k} \quad (5.2)$$

See (2.38), (3.20), §4.2 and (4.47). Thus

$$d\widehat{p_{(i)}} = \sum_{j \in I_{[m]}} r_j^{m-k} dr_j \quad (5.3)$$

and hence

$$\xi_{\widehat{p_{(i)}}} = \sum_{j \in I_{[m]}} r_j^{m-k} \xi_{r_j} \quad (5.4)$$

(see (1.14)).

Now let $z \in M_\Omega(n, \mathfrak{e})$ and let $\nu = \kappa_n(z)$ (see (4.9)) so that $z \in M_\nu(n, \mathfrak{e})$. Let $x = \pi_n(z)$ so that $x \in M_\Omega(n)$. Since the numbers (the eigenvalues of x_m) $r_j(z), j \in I_{[m]}$, are distinct the van der Monde $m \times m$ matrix

$$C_{k\ell} = r_j(z)^{m-k} \quad (5.5)$$

where

$$j = d(m-1) + \ell \quad (5.6)$$

is invertible.

Remark 5.1. In the notation of (4.53) note that $r_j(z) = \nu_{\ell m}$.

If $m = n$ then ξ_{p_i} and $\xi_{p_{(i)}}$ vanish by (2.7) and the argument which implies (2.7). Henceforth assume $m \in I_{n-1}$. Recalling the definition of the m -dimensional commutative Lie algebra $\mathfrak{a}(m)$ of vector fields on $M(n)$ (see §3.1 and (3.20)) there exists a unique basis $\eta_{j\nu}, j \in I_{[m]}$, of $\mathfrak{a}(m)$ such that, for $i \in I_{[m]}$, and k related to i by (5.1),

$$\xi_{p_{(i)}} = \sum_{j \in I_{[m]}} r_j^{m-k}(z) \eta_{j\nu} \quad (5.7)$$

But then (4.46), (5.4), (5.7) and the invertibility of the van der Monde matrix $C_{k\ell}$ implies

Proposition 5.2. *Let $\nu \in \mathfrak{e}_{\Omega(n)}$ and $m \in I_{n-1}$. Then*

$$(\pi_n)_*(\xi_{r_j}|M_\nu(n, \mathfrak{e})) = \eta_{j\nu}|M_c(n) \quad (5.8)$$

where $c = \beta(\nu)$ (see (4.22)) for all $j \in I_{[m]}$.

We recall in §2.4 that $Z_{x,m}$ is the commutative (associative) algebra of $M(m)$ generated by x_m . Here $x = \pi_n(z)$ so that x_m is regular semisimple and hence $\dim Z_{x,m} = m$. We recall (see §3.1) that $G_{x,m}$ is the (algebraic) subgroup of $Gl(n)$ corresponding to $Z_{x,m}$ when $Z_{x,m}$ is regarded as a Lie subalgebra of $M(n)$. We recall also that $A(m)$ is a simply connected group corresponding to $\mathfrak{a}(m)$ and $\mathfrak{a}(m)$ integrates to an action of $A(m)$ on $M(n)$ (see Theorem 3.3). Next we recall (see (3.6)) that $\rho_{x,m}$ is the homomorphism of $A(m)$ into $G_{x,m}$ whose differential is given by

$$(\rho_{x,m})_*(\xi_{p_{(i)}}) = -(x_m)^{m-k} \quad (5.9)$$

when $i \in I_{[m]}$ has the form (5.1). But then applying $(\rho_{x,m})_*$ to (5.7) one has

$$-(x_m)^{m-k} = \sum_{j \in I_{[m]}} r_j^{m-k}(z) (\rho_{x,m})_*(\eta_j \nu) \quad (5.10)$$

On the other hand put

$$h_{\nu,m} = \text{diag}(\nu_{1m}, \dots, \nu_{mm})$$

recalling (4.53), so that in the notation of §4.1

$$h_{\nu,m} \in \mathfrak{e}(m) \quad (5.11)$$

Note that for the matrix units $e_{\ell\ell}$, $\ell \in I_m$, one has

$$h_{\nu,m} = \sum_{\ell \in I_m} \nu_{\ell m} e_{\ell\ell} \quad (5.12)$$

Now let $g_z \in Gl(m)$ be such that

$$g_z h_{\nu,m} g_z^{-1} = x_m \quad (5.13)$$

(see Remark 5.1). Using the notation of §4.1 note then that

$$g_z \mathfrak{d}(m) g_z^{-1} = Z_{x,m} \quad (5.14)$$

Remark 5.3. Note that g_z is unique in $Gl(m)$ modulo the maximal diagonal torus $Diag(m)$ (see §3.4).

For $j \in I_{[m]}$ let $\varepsilon_{z,j} \in Z_{x,m}$ be the idempotent in $Z_{x,m}$ defined by putting

$$\varepsilon_{z,j} = g_z e_{\ell\ell} g_z^{-1} \quad (5.15)$$

where ℓ is defined by (5.6). Thus (5.12) and (5.13) imply that

$$x_m = \sum_{\ell \in I_m} \nu_{\ell m} \varepsilon_{z,d(m-1)+\ell} \quad (5.16)$$

But the $\varepsilon_{z,d(m-1)+\ell}$, $\ell \in I_m$, are orthogonal idempotents by (5.15). Hence, by Remark 5.1,

$$-(x_m)^{m-k} = - \sum_{\ell \in I_m} \nu_{\ell m}^{m-k} \varepsilon_{z,d(m-1)+\ell} \quad (5.17)$$

Proposition 5.4. *Let $z \in M_\Omega(n, \mathfrak{e})$ and let $\nu = \kappa(z)$ (see (4.9)). Let $x = \pi_n(z)$ (see (4.8)) and let $m \in I_n$. Let $\eta_{j,\nu} \in \mathfrak{a}(m)$, $j \in I_{[m]}$, be the basis of $\mathfrak{a}(m)$ defined by (5.7) so that one has (5.8). Let $(\rho_{x,m})_* : \mathfrak{a}(m) \rightarrow Z_{x,m}$ be the Lie algebra homomorphism defined as in (3.6) and (3.7). Let $\varepsilon_{z,j}$, $j \in I_{[m]}$, be the orthogonal idempotents in $Z_{x,m}$ defined by (5.15). Then*

$$(\rho_{x,m})_*(\eta_{j,\nu}) = -\varepsilon_{z,j} \quad (5.18)$$

for any $j \in I_{[m]}$.

Proof. One has $\nu_{\ell m}^{m-k} = r_j^{m-k}(z)$ by Remark 5.1. But then (5.18) follows from the equality of the right hand sides of (5.10) and (5.17), recalling the invertibility of the van der Monde matrix (5.5). QED

5.2. Retain the notation of Proposition 5.4. For any $\zeta \in \mathbb{C}^\times$ and $i \in I_n$ let $\delta_i(\zeta) \in \text{Diag}(n)$ (see §3.4) be the invertible $n \times n$ diagonal matrix such that $\alpha_{j,j}(\delta_i(\zeta)) = 1$ if $j \neq i$ and $\alpha_{i,i}(\delta_i(\zeta)) = \zeta$, using the notation of (1.2). One notes that if $\ell \in I_m$ then $\delta_\ell(\zeta) \in \text{Diag}(m)$. Also, using the relation between j and ℓ given by (5.6), one has $\gamma_{z,j}(\zeta) \in G_{x,m}$, by (5.14), where we put

$$\gamma_{z,j}(\zeta) = g_z \delta_\ell(\zeta) g_z^{-1} \quad (5.19)$$

Let $q \in \mathbb{C}$. One notes that

$$\exp q e_{\ell\ell} = \delta_\ell(e^q) \quad (5.20)$$

and hence

$$\exp q \varepsilon_{z,j} = \gamma_{z,j}(e^q) \quad (5.21)$$

Multiplying (5.18) by q and exponentiating (where $\exp q \eta_{j,\nu} \in A(m)$) it follows then from (3.6) that

$$\rho_{x,m}(\exp q \eta_{j,\nu}) = \gamma_{z,j}(e^{-q}) \quad (5.22)$$

We can now describe the flow generated by ξ_{τ_j} (see (4.47) and §4.2) on $M_\Omega(n, \mathfrak{e})$ for any $j \in I_{d(n-1)}$ (see Theorem 4.27).

Theorem 5.5. *Let $j \in I_{d(n-1)}$ and let $m \in I_{n-1}$ be such that $j \in I_{[m]} = I_{d(m)} - I_{d(m-1)}$. Let $z \in M_\Omega(n, \mathfrak{e})$ (see (4.7)) and let $x \in M_\Omega(n)$, $\nu \in \mathfrak{e}_{\Omega(n)}$ be such that $x = \pi_n(z)$ and $\nu = \kappa_n(z)$. See (4.8) and (4.9). Let $q \in \mathbb{C}$. Then $(\nu, \text{Ad}(\gamma_{z,j}(e^{-q}))(x)) \in M_\nu(n, \mathfrak{e})$ (see (4.7) and (4.21)) and*

$$(\exp q \xi_{\tau_j}) \cdot z = (\nu, \text{Ad}(\gamma_{z,j}(e^{-q}))(x)) \quad (5.23)$$

where $\gamma_{zj}(e^{-q}) \in G_{x,m}$ is defined by (5.21).

Proof. Let $c = \beta(\nu)$ (see (4.22)) so that $c \in \Omega(n)$ and, by Proposition 4.20, the restriction of π_n to $M_\nu(n, \mathfrak{e})$ defines an algebraic isomorphism $M_\nu(n, \mathfrak{e}) \rightarrow M_c(n)$. But then by (5.8) one has

$$\pi_n((\exp q \xi_{r_j}) \cdot z) = (\exp q \eta_{j,\nu}) \cdot x \quad (5.24)$$

(see Theorems 3.23 and 4.27). But $(\exp q \eta_{j,\nu}) \in A(m)$ since by definition $\eta_{j,\nu} \in \mathfrak{a}(m)$. Hence, by (5.22) and Theorem 3.3, one has $(\exp q \eta_{j,\nu}) \cdot x = \text{Ad}(\gamma_{zj}(e^{-q}))(x)$. But then (5.23) follows from (4.7). QED

5.3. We continue with the notation of §5.1 and §5.2.

Proposition 5.6. *Let $j \in I_{d(n-1)}$. Then the isomorphism $\exp q \xi_{r_j}$ of $M_\Omega(n, \mathfrak{e})$ reduces to the identity if $q \in 2\pi i \mathbb{Z}$.*

Proof. This is immediate from (5.23) since $\gamma_{zj}(e^{-q})$ is the identity matrix of $M(n)$, by (5.15) and (5.21), if $q \in 2\pi i \mathbb{Z}$. QED

Let $\mathfrak{r}$ be the commutative $d(n-1)$ dimensional Lie algebra of vector fields on $M_\Omega(n, \mathfrak{e})$ with basis ξ_{r_j} , $j \in I_{d(n-1)}$ (see (4.57) and (4.59)). Let $j \in I_{d(n-1)}$ and let $A_{\mathfrak{r},j}$ be a 1-dimensional complex torus with a global coordinate ζ_j defining an algebraic group isomorphism

$$\zeta_j : A_{\mathfrak{r},j} \rightarrow \mathbb{C}^\times \quad (5.25)$$

By Proposition 5.6 (and abuse of notation) we can regard

$$\mathbb{C} \xi_{r_j} = \text{Lie } A_{\mathfrak{r},j} \quad (5.26)$$

and simultaneously have $A_{\mathfrak{r},j}$ operate on $M_\Omega(n, \mathfrak{e})$, as an integration of the vector field ξ_{r_j} , in such a fashion that if $b \in A_{\mathfrak{r},j}$ then $b = \exp q \xi_{r_j}$ if

$$\zeta_j(b) = e^q \quad (5.27)$$

It is very easy to prove that the action of $A_{\mathfrak{r},j}$ is analytic but what is much more important for us is to prove that this action is that of an algebraic group, operating algebraically on an affine algebraic variety.

Theorem 5.7. *Let $j \in I_{d(n-1)}$. Then the map*

$$A_{\mathfrak{r},j} \times M_\Omega(n, \mathfrak{e}) \rightarrow M_\Omega(n, \mathfrak{e}), \quad (b, z) \mapsto b \cdot z \quad (5.28)$$

is a (algebraic) morphism.

Proof. Let $b \in A_{\mathfrak{r},j}$ and let $z \in M_{\Omega}(n, \mathfrak{e})$. Let $\zeta = \zeta_j(b)$ so that $\zeta \in \mathbb{C}^{\times}$. Recalling (4.7) write $z = (\nu, x)$ where $x \in M_{\Omega}(n)$ and $\nu \in \mathfrak{e}_{\Omega(n)}$. Let $m \in I_{n-1}$ be such that $j \in I_{[m]} = I_{d(m)} - I_{d(m-1)}$. Then, by (5.23) and (5.27),

$$b \cdot z = (\nu, Ad(\gamma_{z,j}(\zeta^{-1}))(x)) \quad (5.29)$$

Since $\nu = \kappa_n(z)$ and $x = \pi_n(z)$ (see (4.8) and (4.9)) and both (4.8) and (4.9) are morphisms it suffices by (5.29) to prove that the map

$$A_{\mathfrak{r},j} \times M_{\Omega}(n, \mathfrak{e}) \rightarrow G_{x,m}, \quad (b, z) \mapsto \gamma_{z,j}(\zeta^{-1}) \quad (5.30)$$

is a morphism. Now $\ell \in I_m$ in (5.19) is defined so that $j = d(m-1) + \ell$. For any $g \in Gl(m)$ let $[g] \in Gl(m)/Diag(m)$ be the left coset defined by g (see §3.4). Of course $Gl(m)/Diag(m)$ is an affine algebraic homogeneous space. Recalling (5.19) to prove (5.30) is a morphism it clearly suffices to show that

$$A_{\mathfrak{r},j} \times M_{\Omega}(n, \mathfrak{e}) \rightarrow Gl(m)/Diag(m), \quad (b, z) \mapsto [g_z] \quad (5.31)$$

is a morphism.

Let $E(m)$ be the set of all regular semisimple elements in $M(m)$ so that $E(m)$ has the structure of a Zariski open (and hence nonsingular) affine subvariety of $M(n)$. Then, recalling (4.1), the map

$$Gl(m) \times \mathfrak{e}(m) \rightarrow E(m), \quad (g, \mu) \mapsto g \mu g^{-1} \quad (5.32)$$

is a surjective morphism. But now if $g \in Gl(m)$ and $\mu \in \mathfrak{e}(m)$ then $[g] \cdot \mu \in E(m)$ is well defined by putting $[g] \cdot \mu = g \mu g^{-1}$. Clearly

$$(Gl(m)/Diag(m)) \times \mathfrak{e}(m) \rightarrow E(m), \quad ([g], \mu) \mapsto [g] \cdot \mu \quad (5.33)$$

is then also a surjective morphism. Now let

$$E(m, \mathfrak{e}) = \{(\mu, y) \in \mathfrak{e}(m) \times E(m) \mid \mu \text{ is } Gl(m)\text{-conjugate to } y\} \quad (5.34)$$

so that $E(m, \mathfrak{e})$ is a Zariski closed subset of $\mathfrak{e}(m) \times E(m)$. The argument establishing the dimension and nonsingularity in Theorem 4.9 (especially using the independence of the differentials dp_i , $i \in I_{[m]}$, at all points in $E(m)$) can obviously be modified to apply here and prove that

$$E(m, \mathfrak{e}) \text{ is a nonsingular } m^2\text{-dimensional Zariski closed subset of } \mathfrak{e}(m) \times E(m) \quad (5.35)$$

But now (5.33) may be augmented to define the map

$$(Gl(m)/Diag(m)) \times \mathfrak{e}(m) \rightarrow E(m, \mathfrak{e}), \quad ([g], \mu) \mapsto (\mu, [g] \cdot \mu) \quad (5.36)$$

But (5.33) readily implies that (5.36) is a surjective morphism so that, for one thing, $E(m, \mathfrak{e})$ is irreducible. Hence $E(m, \mathfrak{e})$ is a nonsingular variety. But (5.36) is obviously bijective and hence birational. But then (5.36) is an algebraic isomorphism. Let

$$E(m, \mathfrak{e}) \rightarrow (Gl(m)/Diag(m)) \times \mathfrak{e}(m) \quad (5.37)$$

be the inverse isomorphism. Projecting on the first factor defines a morphism $\sigma : E(m, \mathfrak{e}) \rightarrow Gl(m)/Diag(m)$ where, for $g \in Gl(m)$ and $\mu \in \mathfrak{e}(m)$,

$$\sigma((\mu, [g] \cdot \mu)) = [g] \quad (5.38)$$

But now since (4.8) and (4.9) are morphisms it follows that $\tau : A_{\mathfrak{r},j} \times M_{\Omega}(n, \mathfrak{e}) \rightarrow E(m, \mathfrak{e})$ is a morphism where, using the notation of (4.3), $\tau((b, z)) = (\nu(m), x_m)$. But

$$\sigma \circ \tau((b, z)) = [g_z] \quad (5.39)$$

by (5.13), since clearly $h_{\nu,m} = \nu(m)$. See (4.3) and (5.12). This proves that (5.31) is a morphism. QED

5.4. Let $m \in I_{n-1}$ and let $\mathfrak{r}(m)$ be the span of the vector fields ξ_{r_i} , $i \in I_{[m]}$, and so that, as defined in §5.3,

$$\mathfrak{r} = \mathfrak{r}(1) \oplus \cdots \oplus \mathfrak{r}(m-1) \quad (5.40)$$

By (4.57) and (4.59), $\mathfrak{r}(m)$ is a commutative Lie algebra of dimension m and as we have already noted $\mathfrak{r}$ is a commutative Lie algebra of dimension $d(n-1)$. Let (see (5.26))

$$\begin{aligned} A_{\mathfrak{r}}(m) &= A_{\mathfrak{r},d(m-1)+1} \times \cdots \times A_{\mathfrak{r},d(m-1)+m} \\ A_{\mathfrak{r}} &= A_{\mathfrak{r}}(1) \times \cdots \times A_{\mathfrak{r}}(n-1) \end{aligned} \quad (5.41)$$

so that as algebraic groups

$$\begin{aligned} A_{\mathfrak{r}}(m) &\cong (\mathbb{C}^\times)^m \\ A_{\mathfrak{r}} &\cong (\mathbb{C}^\times)^{d(n-1)} \end{aligned} \quad (5.42)$$

In addition, by (5.26),

$$\begin{aligned} \mathfrak{r}(m) &= Lie A_{\mathfrak{r}}(m) \\ \mathfrak{r} &= Lie A_{\mathfrak{r}} \end{aligned} \quad (5.43)$$

As an immediate consequence of Theorem 5.7 and commutativity one has

Theorem 5.8. *Let $m \in I_{n-1}$. Then the Lie algebras $\mathfrak{r}(m)$ and $\mathfrak{r}$, respectively, integrate to an algebraic action of $A_{\mathfrak{r}}(m)$ and $A_{\mathfrak{r}}$ on $M_{\Omega}(n, \mathfrak{e})$.*

The following result is a refinement of Theorem 4.25. We are now dealing with the “eigenvalue” vector fields ξ_{r_i} themselves on $M_{\Omega}(n, \mathfrak{e})$ rather than the more crude “eigenvalue symmetric function” vector fields $\widehat{\xi_{p(i)}}$.

Theorem 5.9. *Let $\nu \in \mathfrak{e}_{\Omega(n)}$. Then $M_{\nu}(n, \mathfrak{e})$ is stable under the algebraic group $A_{\mathfrak{r}}$. Furthermore $A_{\mathfrak{r}}$ operates simply and transitively on $M_{\nu}(n, \mathfrak{e})$. In particular the disjoint union (4.21) is the $A_{\mathfrak{r}}$ -orbit decomposition of $M_{\Omega}(n, \mathfrak{e})$.*

Proof. By Theorem 4.22 and Theorem 4.27 one has

$$\mathfrak{r}|M_{\nu}(n, \mathfrak{e}) = \widehat{\mathfrak{a}}|M_{\nu}(n, \mathfrak{e}) \quad (5.44)$$

Hence

$$A_{\mathfrak{r}}|M_{\nu}(n, \mathfrak{e}) = \widehat{A}|M_{\nu}(n, \mathfrak{e}) \quad (5.45)$$

But then $M_{\nu}(n, \mathfrak{e})$ is stable under $A_{\mathfrak{r}}$ and $A_{\mathfrak{r}}$ operates transitively on $M_{\nu}(n, \mathfrak{e})$ by Theorem 4.25. The only question concerns the simplicity of this action.

Let $b \in A_{\mathfrak{r}}$. By definition there exists $q_j \in \mathbb{C}$, $j \in I_{d(n-1)}$, such that

$$b = \exp q_1 \xi_{r_1} \cdots \exp q_{d(n-1)} \xi_{r_{d(n-1)}} \quad (5.46)$$

But then if $c = \beta(\nu)$ (see (4.22)) it follows from Proposition 5.2 that there exists $\eta_{j\nu} \in \mathfrak{a}$, $j \in I_{d(n-1)}$, such that if $a \in A$ is defined by putting

$$a = \exp q_1 \eta_{1,\nu} \cdots \exp q_{d(n-1)} \eta_{d(n-1),\nu} \quad (5.47)$$

then

$$b|M_{\nu}(n, \mathfrak{e}) = \widehat{a}|M_{\nu}(n, \mathfrak{e}) \quad (5.48)$$

recalling Theorem 4.25. In addition Proposition 5.2 implies that $a = a(1) \cdots a(n-1)$ where, for $m \in I_{n-1}$, $a(m) \in A(m)$ is given by

$$a(m) = \prod_{j \in I_{[m]}} \exp q_j \eta_{j,\nu} \quad (5.49)$$

Now assume that $b|M_{\nu}(n, \mathfrak{e})$ has a fixed point. But then by the commutativity of $A_{\mathfrak{r}}$ and the transitivity of $A_{\mathfrak{r}}$ on $M_{\nu}(n, \mathfrak{e})$ it follows that $b|M_{\nu}(n, \mathfrak{e})$ reduces to the identity. We must prove

$$q_j \in 2\pi i \mathbb{Z}, \quad \forall j \in I_{d(n-1)} \quad (5.50)$$

by (5.27) and (5.41). But now $a|M_c(n)$ reduces to the identity by (4.66) and (5.48). See Proposition 4.20. But then $a \in D_c$ (see §3.64). Hence $a(m) \in D_c(m)$ for any $m \in I_{n-1}$ by Theorem 3.28. But Theorem 3.28 also asserts that if $z \in M_\nu(n, \mathfrak{e})$ and $x = \pi_n(z)$ then also $a(m) \in \text{Ker } \rho_{x,m}$. But by (5.15) and (5.18) (see also (5.19) and (5.22)) this implies (5.50) since $m \in I_{n-1}$ is arbitrary. QED

5.5. In the introduction, §0, we defined $\mathfrak{b}_e \subset M(n)$. In common parlance (for some people) $\mathfrak{b}_e$ is the space of all $n \times n$ Hessenberg matrices. In §4.3 we defined $\mathfrak{b}_{e, \Omega(n)}$ to be the intersection $M_\Omega(n) \cap \mathfrak{b}_e$ so that $\mathfrak{b}_{e, \Omega(n)}$ is a Zariski open subvariety of $\mathfrak{b}_e$ (and hence a nonsingular variety) and a closed subvariety of $M_\Omega(n)$. We also defined $M_\Omega(n, \mathfrak{e}, \mathfrak{b}) = \pi_n^{-1}(\mathfrak{b}_{e, \Omega(n)})$ (see (4.8) and §4.3) so that $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ is a Zariski closed subset of $M_\Omega(n, \mathfrak{e})$. Sharpening Theorem 4.4 we shall need

Theorem 5.10. *The restriction (see (4.9))*

$$\kappa_n : M_\Omega(n, \mathfrak{e}, \mathfrak{b}) \rightarrow \mathfrak{e}_{\Omega(n)} \quad (5.51)$$

is an algebraic isomorphism so that $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ is a closed nonsingular subvariety of $M_\Omega(n, \mathfrak{e})$.

Proof. Clearly (5.51) is a morphism since it the restriction of (4.9) to a Zariski closed subset of $M_\Omega(n, \mathfrak{e})$. On the other hand it is bijective by Theorem 4.4. However the inverse of (5.51) is a morphism. See (4.20). QED

An easy consequence of Theorem 5.10 is

Theorem 5.11. *The image of the map*

$$M_\Omega(n, \mathfrak{e}, \mathfrak{b}) \rightarrow \mathbb{C}^{d(n)}, \quad y \mapsto (r_1(y), \dots, r_{d(n)}(y)) \quad (5.52)$$

is a Zariski open set in $\mathbb{C}^{d(n)}$ of the form $(\mathbb{C}^{d(n)})_q$ where q is a nonzero polynomial on $\mathbb{C}^{d(n)}$ and (5.52) is an algebraic isomorphism of $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ with $(\mathbb{C}^{d(n)})_q$.

Proof. Recalling the definition of $\mathfrak{e}_{\Omega(n)}$ in §4.1, Theorem 5.11 follows immediately from Theorem 5.10 and the definition of r_i in (4.47) and $\rho_{k,m}$ in §4.2. QED

On the other hand we establish the following product structure for $M_\Omega(n, \mathfrak{e})$.

Theorem 5.12. *The map*

$$A_\tau \times M_\Omega(n, \mathfrak{e}, \mathfrak{b}) \rightarrow M_\Omega(n, \mathfrak{e}), \quad (b, y) \mapsto b \cdot y \quad (5.53)$$

is an algebraic isomorphism.

Proof. The map (5.53) is bijective by (4.21), (4.25) and Theorem 5.9. But then (5.53) is a bijective morphism of nonsingular algebraic varieties by Theorem 5.8 and Theorem 5.10. Hence (5.53) is an algebraic isomorphism. QED

By definition (see (5.41)) any element $b \in A_{\mathfrak{r}}$ can be uniquely written

$$b = (b_1, \dots, b_{d(n-1)}) \quad (5.54)$$

where $b_j \in A_{\mathfrak{r},j}$. We now extend the domain of the function ζ_j on $A_{\mathfrak{r},j}$ to all of $A_{\mathfrak{r}}$ so that, in the notation of (5.54),

$$\zeta_j(b) = \zeta_j(b_j) \quad (5.55)$$

Thus $\zeta_j \in \mathcal{O}(A_{\mathfrak{r}})$ and the map

$$A_{\mathfrak{r}} \rightarrow (\mathbb{C}^\times)^{d(n-1)}, \quad b \mapsto (\zeta_1(b), \dots, \zeta_{d(n-1)}(b)) \quad (5.56)$$

is an isomorphism of algebraic groups. For $i \in d(n-1)$ let λ_{r_i} be the left invariant vector field on $A_{\mathfrak{r}}$ whose value at the identity of $A_{\mathfrak{r}}$ corresponds to (abuse of notation) ξ_{r_i} . Thus, by (5.27), in the coordinates ζ_j of $A_{\mathfrak{r}}$, one has

$$\lambda_{r_i} = \zeta_i \frac{\partial}{\partial \zeta_i}$$

and hence

$$\lambda_{r_i} \zeta_j = \delta_{ij} \zeta_i \quad (5.57)$$

But now, by Theorem 5.12, every $z \in M_{\Omega}(n, \mathfrak{e})$ can be uniquely written $z = b \cdot y$ where $b \in A_{\mathfrak{r}}$ and $y \in M_{\Omega}(n, \mathfrak{e}, \mathfrak{b})$. Hence, by Theorem 5.12, one has a well-defined function $s_j \in \mathcal{O}(M_{\Omega}(n, \mathfrak{e}))$, $j \in I_{d(n-1)}$, where

$$s_j(z) = \zeta_j(b^{-1}) \quad (5.58)$$

Furthermore Theorems 5.8, 5.9, 5.12 and (5.55) also clearly imply

Theorem 5.13. *Let $\nu \in \mathfrak{e}_{\Omega(n)}$. Then the map*

$$M_{\nu}(n, \mathfrak{e}) \rightarrow (\mathbb{C}^\times)^{d(n-1)}, \quad z \mapsto (s_1(z), \dots, s_{d(n-1)}(z))$$

is an algebraic isomorphism.

The action of $A_{\mathfrak{r}}$ on $M_{\Omega}(n, \mathfrak{e})$ of course introduces, contragrediently, an action of $A_{\mathfrak{r}}$ on $\mathcal{O}(M_{\Omega}(n, \mathfrak{e}))$ so that if $b \in A_{\mathfrak{r}}$, $f \in \mathcal{O}(M_{\Omega}(n, \mathfrak{e}))$ and $z \in M_{\Omega}(n, \mathfrak{e})$ then $(b \cdot f)(z) = f(b^{-1} \cdot z)$. It is immediate from (5.58) that

$$b \cdot s_j = \zeta_j(b) s_j, \quad j \in I_{d(n-1)} \quad (5.59)$$

But then with regard to Poisson bracket on $M_{\Omega}(n, \mathfrak{e})$, at this stage, we can say, for $i \in I_{d(n)}$ and $j \in I_{d(n-1)}$,

$$[r_i, s_j] = \delta_{ij} s_j \quad (5.60)$$

since, by differentiating (5.59) and applying (5.57) clearly,

$$\xi_{r_i} s_j = \delta_{ij} s_j \quad (5.61)$$

recalling (4.59).

Combining Theorems 5.11, 5.12, 5.13 and (5.61) one sees that the s_i, r_j , $i \in I_{d(n-1)}$, $j \in I_{d(n)}$, form a system of uniformizing parameters on $M_{\Omega}(n, \mathfrak{e})$.

Theorem 5.14. *For any $z \in M_{\Omega}(n, \mathfrak{e})$ the n^2 differentials*

$$(dr_i)_z, (ds_j)_z, \quad i \in I_{d(n)}, j \in I_{d(n-1)}, \quad \text{are a basis of the cotangent space } T_z^*(M_{\Omega}(n, \mathfrak{e})) \quad (5.62)$$

Furthermore the image of the map

$$M_{\Omega}(n, \mathfrak{e}) \rightarrow \mathbb{C}^{n^2}, \quad z \mapsto (r_1(z), \dots, r_{d(n)}(z), s_1(z), \dots, s_{d(n-1)}(z)) \quad (5.63)$$

is a Zariski open set Y in $\mathbb{C}^{n^2}$ and (5.63) is an algebraic isomorphism of $M_{\Omega}(n, \mathfrak{e})$ with Y .

Proof. Let $z \in M_{\Omega}(n, \mathfrak{e})$. The $(dr_i)_z$, $i \in I_{d(n)}$, are linearly independent by Proposition 4.18 (double prime statement). On the other hand $(ds_j)_z$, $i \in I_{d(n)}$, $j \in I_{d(n-1)}$, are clearly linearly independent by (5.61). But (5.61) together with (4.57) implies that $(dr_i)_z$ are independent of the $(ds_j)_z$. By dimension this proves (5.62). The image of (5.63) is the just the product of the image of (5.52) and (5.59) by Theorem 5.12. Here we are using the constancy of the r_i on $M_{\nu}(n, \mathfrak{e})$ (see (4.53)). But then the image Y of (5.63) is Zariski open in $\mathbb{C}^{n^2}$ by Theorems 5.11 and 5.13. But the morphism (5.63) is bijective by Theorems 5.11, 5.12 and 5.13. Since both Y and $M_{\Omega}(n, \mathfrak{e})$ are nonsingular varieties it follows (a version of Zariski's Main Theorem) that (5.63) is an algebraic isomorphism. QED

5.6. It is our main objective now to prove the s_i , $i \in I_{d(n-1)}$, Poisson commute among themselves. Localizing this problem, letting $z \in M_\Omega(n, \mathfrak{e})$ and $i, j \in I_{d(n-1)}$ it is enough to show that

$$[s_i, s_j](z) = 0 \quad (5.64)$$

Let $\nu_o = \kappa_n(z)$ so that $z \in M_{\nu_o}(n, \mathfrak{e})$. By (4.25) there exists a unique element z_o in $M_\Omega(n, \mathfrak{e}, \mathfrak{b}) \cap M_{\nu_o}(n, \mathfrak{e})$. Let $x_o = \pi_n(z_o)$ (see (4.8)). Recalling the definition of $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ and $\mathfrak{b}_{e, \Omega(n)}$ in §4.3 it follows that $x_o \in \mathfrak{b}_{e, \Omega(n)}$ and, by Proposition 4.14, π_n defines $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ as an analytic covering space of $\mathfrak{b}_{e, \Omega(n)}$ (see also Theorem 5.10). Thus there exists an open connected (in the Euclidean sense) neighborhood V_{x_o} of x_o in $\mathfrak{b}_{e, \Omega(n)}$ such that V_{x_o} is evenly covered by π_n . Let V_{z_o} be the connected component of $\pi_n^{-1}(V_{x_o})$ which contains z_o . Thus V_{z_o} is an open connected neighborhood of z_o in $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ and

$$\pi_n : V_{z_o} \rightarrow V_{x_o} \quad (5.65)$$

is an analytic isomorphism.

Now, recalling the analytic isomorphism (4.13), let $c_o = \Phi_n(x_o)$ (see (2.8)) and let V_{c_o} be the open connected neighborhood of c_o in $\Omega(n)$ defined by putting $V_{c_o} = \Phi_n(V_{x_o})$. Finally let $V_{\nu_o} = \kappa_n(V_{z_o})$ so that (see (5.51)) V_{ν_o} is open connected neighborhood of ν_o in $\mathfrak{e}_{\Omega(n)}$.

Lemma 5.15. *The map*

$$\beta \circ \kappa_n : V_{z_o} \rightarrow V_{c_o} \quad (5.66)$$

is an analytic isomorphism (see (4.15) for the definition of β).

Proof. One readily notes that $\beta \circ \kappa_n$ restricted to V_{z_o} is the same as $\Phi_n \circ \pi_n$ restricted to V_{z_o} (a commutative diagram). But (5.65) is an analytic isomorphism and (4.13) is an analytic isomorphism. QED

Let $x = \pi_n(z)$. Let $W_z = A_{\mathfrak{r}} \cdot V_{z_o}$ and let $W_x = \pi_n(W_z)$.

Proposition 5.16. *W_z is an open connected neighborhood of z in $M_\Omega(n, \mathfrak{e})$. Furthermore*

$$W_z = \sqcup_{\nu \in V_{\nu_o}} M_\nu(n, \mathfrak{e}) \quad (5.67)$$

In addition W_x is a open connected neighborhood of x in $M_\Omega(n)$ and

$$W_x = \sqcup_{c \in V_{c_o}} M_c(n) \quad (5.68)$$

(see §2.2). Finally

$$\pi_n : W_z \rightarrow W_x \quad (5.69)$$

is an analytic isomorphism.

Proof. Since V_{z_o} is open and connected in $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ it follows from Theorem 5.12 that W_z is open and connected in $M_\Omega(n, \mathfrak{e})$. Furthermore (5.67) follows from Theorems 5.9 and 5.12. Also $z \in W_z$ since $\nu_o \in V_{\nu_o}$. Since a covering map is an open map it follows that W_x is an open connected neighborhood of x in $M_\Omega(n)$. Also (5.68) follows from (5.67) and Proposition 4.20. Since (5.69) is both open and continuous, to prove that it is an analytic isomorphism it suffices to prove that it is bijective. For this of course one must see that it is injective. But this clearly follows from Proposition 4.20, Lemma 5.15 and the bijectivity of (5.51). QED

Proposition 5.16 enables us to carry holomorphic functions on W_z to W_x . Using the notation of §1.2, for $i \in I_{d(n)}$, $j \in I_{d(n-1)}$, let $r'_i, s'_j \in \mathcal{H}(W_x)$ be defined so that on W_z , $r'_i \circ \pi_n = r_i$ and $s'_j \circ \pi_n = s_j$. Recalling the definition of the Poisson structure on $M_\Omega(n, \mathfrak{e})$ (see §4.6) it follows that (5.69) is an isomorphism of Poisson manifolds. Hence to prove (5.64) it suffices to prove

$$[s'_i, s'_j](x) = 0 \quad (5.70)$$

Obviously, for $i \in I_{d(n-1)}$,

$$(\pi_n)_*(\xi_{r'_i}|W_z) = \xi_{r'_i} \quad (5.71)$$

If $\mathfrak{r}'$ is the commutative Lie algebra span of $\xi_{r'_i}$, $i \in I_{d(n-1)}$, then, recalling Proposition 4.20, Theorem 5.9, (5.67) and (5.68), $\mathfrak{r}'$ integrates to a group $A_{\mathfrak{r}'}$, which operates on W_x , and which admits an isomorphism

$$A_{\mathfrak{r}} \rightarrow A_{\mathfrak{r}'}, \quad b \mapsto b' \quad (5.72)$$

such that for any $w \in W_z$, and $b \in A_{\mathfrak{r}}$

$$\pi_n(b \cdot w) = b' \cdot \pi_n(w) \quad (5.73)$$

In addition Theorems 5.9 and 5.12 imply

Proposition 5.17. (5.68) is the orbit decomposition of $A_{\mathfrak{r}'}$ on W_x and $A_{\mathfrak{r}'}$ operates simply (as well as transitively) on $M_c(n)$ for every $c \in V_{c_o}$. Furthermore (see (5.65))

$$W_x = \sqcup_{b \in A_{\mathfrak{r}}} b' \cdot V_{x_o} \quad (5.74)$$

5.7. We retain the notation of the previous section and recall some notation from §1.2. If $O \subset M(n)$ is an adjoint orbit of $Gl(n)$ and $y \in O$ then $O_y = O$. If O is an orbit of regular semisimple elements then

$$\dim O = 2d(n-1) \quad (5.75)$$

and if $q_k, k = 1, \dots, n$, is the constant value that the $Gl(n)$ -invariant (see §2.1) $p_{d(n-1)+k}$ takes on O then O is clearly determined by those values. That is,

$$O = \{y \in M(n) \mid p_{d(n-1)+k}(y) = q_k\} \quad (5.76)$$

It follows then from (2.9) that if (see §2.2)

$$\mathbb{C}^{d(n)}(O) = \{c \in \mathbb{C}^{d(n)} \mid c_{d(n-1)+k} = q_k, k \in I_n\}$$

one has

$$O = \sqcup_{c \in \mathbb{C}^{d(n)}(O)} M_c(n) \quad (5.77)$$

Let $\mathcal{S}$ be the set of all $Gl(n)$ -adjoint orbits of regular semisimple elements. Since any $y \in M_\Omega(n)$ (see (2.53)) is regular semisimple one has $O_y \in \mathcal{S}$ for any $y \in W_x$. Let

$$\mathcal{S}(W_x) = \{O \in \mathcal{S} \mid W_x \cap O \neq \emptyset\}$$

and for any $O \in \mathcal{S}(W_x)$ let $V_{c_o}(O) = \mathbb{C}^{d(n)}(O) \cap V_{c_o}$ so that

$$O \cap W_x = \sqcup_{c \in V_{c_o}(O)} M_c(n) \quad (5.78)$$

by (5.68) and (5.77).

Now, recall from §1.2, any adjoint orbit has the structure of a symplectic manifold. If $O \in \mathcal{S}$ then as one knows O is closed in $M(n)$. If $O \in \mathcal{S}(W_x)$ then $O \cap W_x$ is open in O and hence $O \cap W_x$ is a symplectic manifold.

Proposition 5.18. *Let $O \in \mathcal{S}(W_x)$. Then the group $A_{\mathfrak{r}'}$ (see Proposition 5.17) stabilizes $O \cap W_x$ and operates as a group of symplectomorphisms on $O \cap W_x$.*

Proof. For any $i \in I_{d(n-1)}$ the vector field $\xi_{r'_i}|_{O \cap W_x}$ is tangent to $O \cap W_x$ and is a Hamiltonian vector field on $O \cap W_x$ by Proposition 1.3 and especially (1.19). Thus $\mathfrak{r}'|_{O \cap W_x}$ is a Lie algebra of Hamiltonian vector fields on $O \cap W_x$. But, by (5.78) and

Proposition 5.17, $O \cap W_x$ is stabilized by the integrated group $A_{\mathfrak{r}'}$. Hence $A_{\mathfrak{r}'}$ operates as a group of symplectomorphisms of $O \cap W_x$. QED

Assume X is a submanifold of W_x . Let $\mathcal{S}(X) = \{O \in \mathcal{S} \mid O \cap X \neq \emptyset\}$ so that $\mathcal{S}(X) \subset \mathcal{S}(W_x)$. We will say X is Lagrangian in W_x if $\dim X = d(n)$ and $O \cap X$ is a Lagrangian submanifold of the symplectic manifold O , for any $O \in \mathcal{O}(X)$.

Proposition 5.19. *Assume X is a Lagrangian submanifold of W_x . Let $b \in A_{\mathfrak{r}'}$. Then $\mathcal{S}(X) = \mathcal{S}(b' \cdot X)$ and $b' \cdot X$ is again a Lagrangian submanifold of W_x .*

Proof. Obviously $\dim b' \cdot X = d(n)$. Let $O \in \mathcal{S}(X)$. Then of course $O \cap X = (O \cap W_x) \cap X$ and, by definition of being Lagrangian, obviously $(O \cap W_x) \cap X$ is Lagrangian in $O \cap W_x$. But b' operates as a symplectomorphism of $O \cap W_x$ by Proposition 5.18. Thus $b' \cdot ((O \cap W_x) \cap X) = (O \cap W_x) \cap b' \cdot X$ is again Lagrangian in $O \cap W_x$. That is, $O \in \mathcal{S}(b' \cdot X)$ and $b' \cdot (O \cap X) = O \cap b' \cdot X$ is Lagrangian in O . By using b^{-1} one readily reverses the argument to show that if $O \in \mathcal{S}(b' \cdot X)$ then $O \in \mathcal{S}(X)$ and $O \cap b' \cdot X$ is Lagrangian in O . QED

The following result will be seen to be the key point in proving (5.70) and consequently (5.64).

Theorem 5.20. *Retain the notation of (5.65) (or (5.74)). Then V_{x_o} is a Lagrangian submanifold of W_x .*

Proof. We will use results in [K2]. These are stated for complex semisimple Lie groups but their extension to the reductive group $Gl(n)$ is immediate and we will apply the results for that case. Let $N \subset Gl(n)$ be the maximal unipotent subgroup where $Lie N$ is the Lie subalgebra of all strictly upper triangular matrices. Retaining notation in [KW] we have put $\mathfrak{u} = Lie N$. See (3.51)). Let $\mathfrak{s}$ be defined by (1.1.5) in [K2] so that if $\mathfrak{s}_e = -e + \mathfrak{s}$, using the notation of §2.2, then $\mathfrak{s}_e \subset \mathfrak{b}_e$ (see §4.3) and, as asserted by Theorem 1.1 in [K2], (a), $\mathfrak{s}_e$ is a cross-section for the adjoint action of $Gl(n)$ on the set of all regular elements in $M(n)$. On the other hand (b), Theorem 1.2 in [K2] asserts that $\mathfrak{b}_e$ is stable under $Ad N$ and the map

$$N \times \mathfrak{s}_e \rightarrow \mathfrak{b}_e, \quad (u, w) \mapsto Ad u(w) \quad (5.79)$$

is an isomorphism of affine varieties.

To prove that V_{x_o} is Lagrangian in W_x we first observe that $\dim V_{x_o} = d(n)$. This is clear since V_{x_o} is open in $\mathfrak{b}_e$ (see §5.6). Now let $O \in \mathcal{S}(V_{x_o})$. It remains to show

that $O \cap V_{x_o}$ is Lagrangian in O . But now, by (a) above, $O \cap \mathfrak{s}_e$ consists of a single point y_o and hence by (b) one must have

$$O \cap \mathfrak{b}_e = \text{Ad } N(y_o) \quad (5.80)$$

Since O is closed in $M(n)$ this implies that $\text{Ad } N(y_o)$ is closed in $\mathfrak{b}_e$ and one has

$$O \cap V_{x_o} = \text{Ad } N(y_o) \cap V_{x_o} \quad (5.81)$$

since $V_{x_o} \subset \mathfrak{b}_e$. But V_{x_o} is open in $\mathfrak{b}_e$ and hence (5.81) implies that $O \cap V_{x_o}$ is open in $\text{Ad } N(y_o)$. Consequently to prove that $O \cap V_{x_o}$ is Lagrangian in O it suffices to prove that $\text{Ad } N(y_o)$ is Lagrangian in O . But $\dim \text{Ad } N(y_o) = \dim N$ by (5.79), and $\dim N = d(n-1)$ which is half the dimension of O . On the other hand if $y \in \text{Ad } N(y_o)$ and $u, v \in \mathfrak{u}$ (see (3.51)) we must show that $\omega_y(\eta^u, \eta^v) = 0$ (see §1.2 and more specifically (1.10)). But $\omega_y(\eta^u, \eta^v) = B(y, [u, v])$ by (1.10). But $B(y, [u, v]) = 0$ since $y \in \mathfrak{b}_e$ and one notes that $\mathfrak{b}_e$ is B -orthogonal to $[\mathfrak{u}, \mathfrak{u}]$. QED

By Theorem 5.12 one has the following disjoint union

$$M_\Omega(n, \mathfrak{e}) = \sqcup_{b \in A_\tau} b \cdot M_\Omega(n, \mathfrak{e}, \mathfrak{b}) \quad (5.82)$$

On the other hand, we note that the components in (5.82) are level sets of the functions s_j , $j \in I_{d(n-1)}$. Indeed for any $\tau \in (\mathbb{C}^\times)^{d(n-1)}$ let $\tau_i \in \mathbb{C}^\times$, $i \in I_{d(n-1)}$, be defined so that

$$\tau = (\tau_1, \dots, \tau_{d(n-1)}) \quad (5.83)$$

and for $\tau \in (\mathbb{C}^\times)^{d(n-1)}$, let

$$M_\Omega(n, \mathfrak{e}, \tau) = \{y \in M_\Omega(n, \mathfrak{e}) \mid s_i(y) = \tau_i, \forall i \in I_{d(n-1)}\} \quad (5.84)$$

so that

$$M_\Omega(n, \mathfrak{e}) = \sqcup_{\tau \in (\mathbb{C}^\times)^{d(n-1)}} M_\Omega(n, \mathfrak{e}, \tau) \quad (5.85)$$

The following proposition is an immediate consequence of (5.58).

Proposition 5.21. *The partitions (5.74) and (5.82) of $M_\Omega(n, \mathfrak{e})$ are the same. That is, for any $b \in A_\tau$,*

$$b \cdot M_\Omega(n, \mathfrak{e}, \mathfrak{b}) = M_\Omega(n, \mathfrak{e}, \tau) \quad (5.86)$$

where for $i \in I_{d(n-1)}$, $\tau_i = \zeta_i(b^{-1})$.

Remark 5.22. Since V_{z_o} is open in $M_\Omega(n, \mathfrak{e}, \tau)$ (see §5.6) note from the definition W_z in §5.6 one has the disjoint union

$$W_z = \sqcup_{b \in A_\tau} b \cdot V_{z_o} \quad (5.87)$$

and hence by (5.86),

$$b \cdot V_{z_o} = \{w \in W_z \mid s_i(w) = \zeta_i(b^{-1}), \forall i \in I_{d(n-1)}\} \quad (5.88)$$

We can now prove one of the main theorems of the paper.

Theorem 5.23. *The uniformizing parameters $r_i, s_j, i \in I_{d(n)}, j \in I_{d(n-1)}$ of $M_\Omega(n, \mathfrak{e})$ (see Theorem 5.14) satisfy the following Poisson commutation relations,*

$$\begin{aligned} (1) \quad & [r_i, r_j] = 0, \quad i, j \in I_{d(n)} \\ (2) \quad & [r_i, s_j] = \delta_{ij} s_j, \quad i \in I_{d(n)}, j \in I_{d(n-1)} \\ (3) \quad & [s_i, s_j] = 0, \quad i, j \in I_{d(n-1)} \end{aligned} \quad (5.89)$$

Proof. (1) and (2) have already been proved. See (4.57) and (5.60). We therefore have only to prove (3). We use the notation of §5.6 where we have to prove (5.64). But, as we have observed, this comes down to proving (5.70). Recalling the isomorphism (5.69) and the definitions of s'_i and b' for $b \in A_\tau$ in §5.6, one has the disjoint union

$$W_x = \sqcup_{b \in A_\tau} b' \cdot V_{x_o} \quad (5.90)$$

and

$$b' \cdot V_{x_o} = \{y \in V_x \mid s_i(y) = \zeta_i(b^{-1}), \forall i \in I_{d(n-1)}\} \quad (5.91)$$

by (5.87) and (5.88).

But now, by (5.90), there exists $b \in A_\tau$ such that $x \in b' \cdot V_{x_o}$. For notational simplicity put $X = b' \cdot V_{x_o}$. But then X is Lagrangian submanifold of V_x by Propositions 5.19 and 5.20. Obviously $O_x \in \mathcal{S}(X)$ (using the notation of Proposition 5.19). Thus $O_x \cap X$ is a Lagrangian submanifold of the symplectic manifold O_x and $x \in O_x \cap X$. Let $v \in T_x(O_x \cap X)$. But since the s'_i are constant on X by (5.91) one has $v s'_i = 0$ for all $i \in I_{d(n-1)}$. On the other hand $(\xi_{s'_i})_x \in T_x(O_x)$ by Proposition 1.3. Thus $\omega_x((\xi_{s'_i})_x, v) = 0$, by (1.17) and (1.19), for all v in the Lagrangian subspace $T_x(O_x \cap X)$ of $T_x(O_x)$. But by the isotropic maximality of $T_x(O_x \cap X)$, with respect to ω_x , one must have $(\xi_{s'_i})_x \in T_x(O_x \cap X)$. But then $(\xi_{s'_i})_x s'_j = 0$ for all $i, j \in I_{d(n-1)}$ since s'_j is constant on $O_x \cap X$. This proves (5.70). QED

5.8. The Poisson bracket in $P(n)$ (see §1.1) or, as denoted in §4.4, $\mathcal{O}(M(n))$, extends in the obvious way to the quotient field, $F(n)$ (see §4.5). Similarly, the Poisson structure in $\mathcal{O}(M_\Omega(n, \mathfrak{e}))$ extends to the function field $F(n, \mathfrak{e})$ (see §4.5) which, we recall, is a Galois extension of $F(n)$. See Proposition 4.12.

The Gelfand-Kirillov theorem is the statement that the quotient division ring of the universal enveloping algebra of $M(n)$ is isomorphic to the quotient division ring of a Weyl algebra over a (central) rational function field. A natural commutative analogue is the statement that $F(n)$, as a Poisson field, is isomorphic to the rational function field of a classical phase space over (a Poisson central) function field. Using the eigenvalue functions, r_i , and the commutative algebraic group A_τ we now find that the statement is explicitly true for the Galois extension $F(n, \mathfrak{e})$ of $F(n)$.

Theorem 5.24. *For $i \in I_{d(n-1)}$ one has $s_i^{-1} \in \mathcal{O}(M_\Omega(n, \mathfrak{e}))$ (see (5.58)) so that (see §4.2 and (4.47)) $r_{(i)} \in \mathcal{O}(M_\Omega(n, \mathfrak{e}))$ where*

$$r_{(i)} = r_i / s_i \quad (5.92)$$

Then $F(n, \mathfrak{e})$ is the rational function field in n^2 variables,

$$F(n, \mathfrak{e}) = \mathbb{C}(r_{(1)}, \dots, r_{(d(n-1))}, s_1, \dots, s_{d(n-1)}, r_{d(n-1)+1}, \dots, r_{d(n-1)+n}) \quad (5.93)$$

Furthermore one has the Poisson commutation relations: $r_{d(n-1)+k}$ Poisson commutes with all every element in $F(n, \mathfrak{e})$ for $k \in I_n$ and, for $i, j \in I_{d(n-1)}$,

$$\begin{aligned} (1) \quad & [r_{(i)}, r_{(j)}] = 0 \\ (2) \quad & [r_{(i)}, s_j] = \delta_{ij} \\ (3) \quad & [s_i, s_j] = 0 \end{aligned} \quad (5.94)$$

Proof. As a function field one has

$$F(n, \mathfrak{e}) = \mathbb{C}(r_1, \dots, r_{d(n-1)}, s_1, \dots, s_{d(n-1)}, r_{d(n-1)+1}, \dots, r_{d(n-1)+n}) \quad (5.95)$$

by Theorem 5.14. For $i \in I_{d(n-1)}$ one has $s_i^{-1} \in \mathcal{O}(M_\Omega(n, \mathfrak{e}))$ since s_i vanishes nowhere on $M_\Omega(n, \mathfrak{e})$ by (5.58). As a function field (5.93) follows immediately from (5.95). The commutation relations (5.94) follow easily from (5.89). QED

Involved in the paper are two groups which operate on $M_\Omega(n, \mathfrak{e})$ and then contragrediently on $\mathcal{O}(M_\Omega(n, \mathfrak{e}))$. In both cases the latter action extends to an action on the field $F(n, \mathfrak{e})$. The first group is the Galois group Σ_n (see §4.5 and Proposition

4.12). It is immediate from Proposition 4.14 that Σ_n preserves Poisson bracket. The second group is the $d(n-1)$ -dimensional complex torus, $A_{\mathfrak{r}}$. Since the action of $A_{\mathfrak{r}}$ is algebraic (see Theorem 5.8) the group $A_{\mathfrak{r}}$ stabilizes $\mathcal{O}(M_{\Omega}(n, \mathfrak{e}))$. In fact since the action is algebraic, as one knows, for any $f \in \mathcal{O}(M_{\Omega}(n, \mathfrak{e}))$,

$$A_{\mathfrak{r}} \cdot f \text{ spans a finite dimensional subspace of } \mathcal{O}(M_{\Omega}(n, \mathfrak{e})) \quad (5.96)$$

In addition, the action of $A_{\mathfrak{r}}$ obviously extend to an action on the field $F(n, \mathfrak{e})$. Furthermore this action also preserves Poisson bracket (an easy consequence of (5.46) and (5.96)). The following theorem explicitly determines the action of the two groups. If $\alpha \in \mathbb{Z}^{d(n-1)}$ and $j \in I_{d(n-1)}$ let $\alpha_j \in \mathbb{Z}$ be such that $\alpha = (\alpha_1, \dots, \alpha_{d(n-1)})$. For $\alpha \in \mathbb{Z}$ let ζ^{α} be the character on the torus $A_{\mathfrak{r}}$ defined by putting (see (5.55) and (5.56))

$$\zeta^{\alpha} = \zeta_1^{\alpha_1} \cdots \zeta_{d(n-1)}^{\alpha_{d(n-1)}}$$

Also let $s^{\alpha} \in F(n, \mathfrak{e})$ be defined by putting

$$s^{\alpha} = s_1^{\alpha_1} \cdots s_{d(n-1)}^{\alpha_{d(n-1)}}$$

Theorem 5.25. *Let $m \in I_n$ and let $j \in I_{[m]} = I_{d(m)} - I_{d(m-1)}$. Then for any $\sigma \in \Sigma_n$ there exists $k \in I_{[m]}$ such that*

$$\sigma \cdot r_j = r_k \quad (5.97)$$

Furthermore if $m \in I_{n-1}$ then $\sigma \cdot s_j = s_k$ and $\sigma \cdot r_{(j)} = r_{(k)}$. In addition Σ_n normalizes the torus $A_{\mathfrak{r}}$. In fact in the preceding notation

$$\sigma A_{\mathfrak{r},j} \sigma^{-1} = A_{\mathfrak{r},k} \quad (5.98)$$

(see §5.3 and (5.41).

Next, for any rational function $f \in \mathbb{C}(r_1, \dots, r_{d(n)})$ and any $\alpha \in \mathbb{Z}^{d(n-1)}$ one has

$$b \cdot (f s^{\alpha}) = \zeta^{\alpha}(b) f s^{\alpha} \quad (5.99)$$

for any $b \in A_{\mathfrak{r}}$.

Proof. (5.97) follows immediately from the definition of the action of Σ_n . See (4.5), (4.6), (4.10), (4.11), the definition of ρ_{i_m} in §4.2 and (4.47). Since σ preserves the Poisson bracket structure in $M_{\Omega}(n, \mathfrak{e})$ it carries the vector field ξ_{r_j} to ξ_{r_k} and, hence transforms the corresponding flows (where, by (4.59), we may assume $m \in I_{n-1}$) so

as to yield (5.98). But σ also stabilizes $M_\Omega(n, \mathfrak{e}, \mathfrak{b})$ since, by definition, (see §5.5) $M_\Omega(n, \mathfrak{e}, \mathfrak{b}) = \pi_n^{-1}(\mathfrak{b}_{e, \Omega(n)})$, But then $\sigma \cdot s_j = s_k$ by (5.58) and (5.98). It is, of course, then immediate that $\sigma \cdot r_{(j)} = r_{(k)}$.

The equation (5.99) follows obviously from the definition of A_τ (see (5.41)) and (5.59). QED

References for Part II

- [B] A. Borel, *Linear Algebraic Groups*, W. A. Benjamin, Inc, 1969
- [GKL1] A. Gerasimov, S. Kharchev and D. Lebedev, Representation Theory and Quantum Inverse Scattering Method: The Open Toda Chain and the Hyperbolic Sutherland Model, IMRN 2004 **17**, 2004, 823–854, arXiv:math. QA/0204206
- [GKL2] A. Gerasimov, S. Kharchev and D. Lebedev, On a class of integrable systems connected with $GL(N, \mathbb{R})$, Int. J. Mod. Phys. A, vol 19, Suppl. 2004, 823–854, arXiv: math. QA/0301025
- [K2] B. Kostant, On Whittaker Vectors and Representation Theory, Inventiones Math. vol 48(1978), 101-184
- [KW] B. Kostant and N. Wallach, Gelfand-Zeitlin theory from the perspective of classical mechanics I, <http://www.arXiv.org/pdf/math.SG/0408342>, to appear in Studies in Lie Theory, A. Joseph Festschrift, Birkhauser, PM series.
- [M1] D. Mumford, *The red book of varieties and schemes*, Lecture Notes in Math., 1358, Springer, 1995
- [M2] D. Mumford, Geometric Invariant Theory, Ergeb. Math., Band 34(1965), Springer-Verlag
- [Sm] L. Smith, Polynomial Invariants of Finite Groups, Research Notes in Math. vol 6(1995), A. K. Peters, Ltd
- [Sp] T. Springer, *Linear Algebraic Groups*, 2nd edition, Prog in Math., vol **9**, Birkhauser Boston, 1998
- [ZS] O. Zariski and P. Samuel, *Commutative Algebra*, Vol 1(1958), Van Nostrand.

Bertram Kostant
 Dept. of Math.
 MIT
 Cambridge, MA 02139
 kostant@math.mit.edu

Nolan Wallach
 Dept. of Math.
 UCSD
 San Diego, CA 92093
 nwallach@ucsd.edu